arXiv:0710.3218v1 [math.RT] 17 Oct 2007

REPRESENTATIONS OF TEMPERLEY–LIEB ALGEBRAS

JOHN ENYANG

ABSTRACT. We define a commuting family of operators $T_0, T_1, \ldots, T_n$ in the Temperley–Lieb algebra $\mathcal{A}_n(x)$ of type A_{n-1}. Using an appropriate analogue to Murphy basis of the Iwahori–Hecke algebra of the symmetric group, we describe the eigenvalues arising from the triangular action of the said operators on the cell modules of $\mathcal{A}_n(x)$. These results are used to provide the Temperley–Lieb algebras of type A_{n-1} with a semi–normal form, together with a branching law, and explicit formulae for associated Gram determinants.

1. The Temperley–Lieb Algebras

Let n be a non–negative integer, x be an indeterminate over $\mathbb{Z}$ and write $R = \mathbb{Z}[x]$. The Temperley–Lieb algebra $\mathcal{A}_n(x)$, defined in [9], is the unital associative R–algebra generated by $e_1, \ldots, e_{n-1}$ which are subject to the defining relations

$$e_i^2 = xe_i, \qquad \text{for } i = 1, \ldots, n-1; \tag{1.1}$$

$$e_i e_{i\pm1} e_i = e_i, \qquad \text{for } i, i \pm 1 = 1, \ldots, n-1; \tag{1.2}$$

$$e_i e_j = e_j e_i, \qquad \text{for } i, j = 1, \ldots, n-1 \text{ and } |i - j| \geq 2. \tag{1.3}$$

By convention, $\mathcal{A}_1(x) = R$ and, for $i = 2, 3, \ldots,$ we regard $\mathcal{A}_i(x)$ as the subalgebra of $\mathcal{A}_{i+1}(x)$ generated by $e_1, \ldots, e_{i-1}$, giving a tower

$$\mathcal{A}_1(x) \subseteq \mathcal{A}_2(x) \subseteq \mathcal{A}_3(x) \subseteq \cdots . \tag{1.4}$$

Using restriction in the tower (1.4), we construct cellular bases, in the sense of [2], for $\mathcal{A}_n(x)$ which are compatible with the action of certain commuting operators in $\mathcal{A}_n(x)$.

2. Murphy Bases for the Temperley–Lieb Algebras

For the purposes of these notes, a *partition* of n is a pair of integers $\lambda = (i, n - 2i)$, where $0 \leq 2i \leq n$. If $\lambda = (i, n - 2i)$ and $\mu = (j, n - 2j)$ are partitions of n, we will write $\lambda \trianglerighteq \mu$ if $i \geq j$, while $\lambda \vartriangleright \mu$ will signify that $\lambda \trianglerighteq \mu$ and $\lambda \neq \mu$. If μ is a partition of $n - 1$ and $\lambda = (i, n - 2i)$ is a partition of n, write $\mu \to \lambda$ if $\mu = (i, n - 2i - 1)$ or $\mu = (i - 1, n - 2i + 1)$. Let λ be a partition of n. Define

$$\mathfrak{T}_n(\lambda) = \{(\lambda^{(0)}, \lambda^{(1)}, \ldots, \lambda^{(n)}) : \lambda^{(0)} = (0, 0),\ \lambda^{(k-1)} \to \lambda^{(k)} \text{ for } k = 1, \ldots, n, \text{ and } \lambda^{(n)} = \lambda\}.$$

If $\mathfrak{s} = (\lambda^{(0)}, \ldots, \lambda^{(n)}) \in \mathfrak{T}_n(\lambda)$, write $\lambda = \mathrm{Shape}(\mathfrak{s})$ and let $\mathfrak{s}|_k = (\lambda^{(0)}, \ldots, \lambda^{(k)})$, for $k = 1, \ldots, n$. We order the elements of $\mathfrak{T}_n(\lambda)$ by writing $\mathfrak{s} \trianglerighteq \mathfrak{t}$ if $\mathrm{Shape}(\mathfrak{s}|_k) \trianglerighteq \mathrm{Shape}(\mathfrak{t}|_k)$

Key words and phrases. Temperley–Lieb algebra; Specht module; cellular algebra; Murphy basis; Jucys–Murphy operators.

for $k = 1, \ldots, n$, and $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$; by $\mathfrak{s} \rhd \mathfrak{t}$ we will mean that $\mathfrak{s} \unrhd \mathfrak{t}$ and $\mathfrak{s} \neq \mathfrak{t}$. If $\mathfrak{t} = (\lambda^{(0)}, \ldots, \lambda^{(n)})$, then $\mathfrak{t}$ may be identified with an up–down tableau:

$$\mathfrak{t} \mapsto (\mathfrak{t}^{(0)}, \ldots, \mathfrak{t}^{(n)}), \quad \text{where } \mathfrak{t}^{(k)} = \underbrace{\square\square\cdots\square}_{(k-2i)\text{ boxes}} \text{ whenever } \lambda^{(k)} = (i, k-2i), \text{ for } k = 0, \ldots, n.$$

In turn, the up–down tableaux correspond to paths in the Bratteli diagram associated with the Temperley–Lieb algebras (*cf.* §2 of [5]).

If f is an integer, $0 \leq f \leq [n/2]$ and $\lambda = (f, n-2f)$, define

$$m_\lambda = e_1 e_3 \cdots e_{2f+1}$$

and let $\mathcal{A}_n^\lambda$ denote the two sided ideal in $\mathcal{A}_n(x)$ generated by m_λ and

$$\check{\mathcal{A}}_n^\lambda = \sum_{\mu \rhd \lambda} \mathcal{A}_n^\mu.$$

If $n = 2k + \delta$, where $\delta \in \{0, 1\}$, then

$$0 \subset \mathcal{A}_n^{(k, n-2k)} \subset \mathcal{A}_n^{(k-1, n-2k+2)} \subset \cdots \subset \mathcal{A}_n^{(0,n)} = \mathcal{A}_n(x)$$

is a filtration by two sided ideals of $\mathcal{A}_n(x)$.

If $i, j = 1, 2, \ldots, n$, define $w_{i,j} \in \mathcal{A}_n$ by

$$w_{i,j} = \begin{cases} e_i e_{i+1} \cdots e_{j-1}, & \text{if } i < j; \\ e_{i-1} e_{i-2} \cdots e_j, & \text{if } j < i; \\ 1, & \text{otherwise.} \end{cases}$$

Now, introduce elements

$$\{v_\mathfrak{t} : \mathfrak{t} \in \mathfrak{T}_n(\lambda), \ \lambda \text{ a partition of } n\},$$

by writing $v_\mathfrak{t} = 1$ if $\mathfrak{t} = ((0,0))$ and, otherwise, if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, where $\lambda = (f, n-2f)$, and $\mathfrak{s} = \mathfrak{t}|_{n-1}$, then

$$v_\mathfrak{t} = \begin{cases} v_\mathfrak{s}, & \text{if } \mathrm{Shape}(\mathfrak{s}) = (f, n-2f-1) \\ w_{2f,n} v_\mathfrak{s}, & \text{if } \mathrm{Shape}(\mathfrak{s}) = (f-1, n-2f+1). \end{cases}$$

Similarly, we define

$$\{v_\mathfrak{t}^* : \mathfrak{t} \in \mathfrak{T}_n(\lambda), \ \lambda \text{ a partition of } n\},$$

by writing $v_\mathfrak{t}^* = 1$ if $\mathfrak{t} = ((0,0))$, and, otherwise, if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, where $\lambda = (f, n-2f)$, and $\mathfrak{s} = \mathfrak{t}|_{n-1}$, then

$$v_\mathfrak{t}^* = \begin{cases} v_\mathfrak{s}^*, & \text{if } \mathrm{Shape}(\mathfrak{s}) = (f, n-2f-1) \\ v_\mathfrak{s}^* w_{n,2f}, & \text{if } \mathrm{Shape}(\mathfrak{s}) = (f-1, n-2f+1). \end{cases}$$

The following are stated for reference in subsequent calculations.

Lemma 2.1. *Suppose that $\lambda = (f, n-2f)$ is a partition of n, with $f > 0$ and $n - 2f \geq 1$. Let $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$ satisfy $\mathfrak{s}|_{n-2} = \mathfrak{t}|_{n-2}$ and $\mathfrak{s} \neq \mathfrak{t}$. Then the condition*

$$\mathrm{Shape}(\mathfrak{s}|_{n-1}) = (f-1, n-2f+1), \qquad \textit{and} \qquad \mathrm{Shape}(\mathfrak{t}|_{n-1}) = (f, n-2f-1)$$

holds if and only if $v_\mathfrak{s} = v_\mathfrak{t} e_{n-1}$.

Proof. Suppose that $\mathfrak{s} \neq \mathfrak{t}$ and $\mathfrak{s}|_{n-2} = \mathfrak{t}|_{n-2}$ and consider the Bratteli diagram fragment

$$\begin{tikzcd}
 & (f, n-2f-1) \arrow[dr] & \\
(f-1, n-2f) \arrow[ur] \arrow[dr] & & (f, n-2f) \\
 & (f-1, n-2f+1) \arrow[ur] \arrow[dr] & \\
 & & (f-1, n-2f+2).
\end{tikzcd}$$

If $\mathfrak{u} = \mathfrak{t}|_{n-2}$, then either,

$$\operatorname{Shape}(\mathfrak{s}|_{n-1}) = (f, n-2f-1) \qquad \text{and} \qquad \operatorname{Shape}(\mathfrak{t}|_{n-1}) = (f-1, n-2f+1),$$

in which case $v_\mathfrak{s} = w_{2f,n-1} v_\mathfrak{u}$ and $v_\mathfrak{t} = w_{2f,n} v_\mathfrak{u}$, or

$$\operatorname{Shape}(\mathfrak{s}|_{n-1}) = (f-1, n-2f+1) \qquad \text{and} \qquad \operatorname{Shape}(\mathfrak{t}|_{n-1}) = (f, n-2f-1),$$

in which case $v_\mathfrak{s} = w_{2f,n} v_\mathfrak{u}$ and $v_\mathfrak{t} = w_{2f,n-1} v_\mathfrak{u}$. Since $w_{2f,n} = w_{2f,n-1} e_{n-1}$, and e_{n-1} commutes with $v_\mathfrak{u}$, the result follows. □

Corollary 2.2. *Let λ be a partition of n and suppose that k is an integer, $1 < k < n$. If $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$ satisfy $\mathfrak{s} \neq \mathfrak{t}$ and* $\operatorname{Shape}(\mathfrak{s}|_i) = \operatorname{Shape}(\mathfrak{t}|_i)$, *for $i \in \{0, 1, \ldots, n\} \setminus \{k-1\}$, then the condition*

$$\operatorname{Shape}(\mathfrak{s}|_{k-1}) = (j-1, k-2j+1), \qquad \textit{and} \qquad \operatorname{Shape}(\mathfrak{t}|_{k-1}) = (j, k-2j-1)$$

holds if and only if $v_\mathfrak{s} = v_\mathfrak{t} e_{k-1}$.

In [2], J. Graham and G. Lehrer have demonstrated that $\mathcal{A}_n(x)$ is cellular, while M. Härterich has provided certain Murphy type bases for generalised Temperley–Lieb algebras in [7]. In order to obtain a triangular action for the commuting family of elements defined in §3, we establish that $\mathcal{A}_n(x)$ has a cellular basis as described in the next lemma (*cf.* Example 2.1 below).

Lemma 2.3. *The algebra $\mathcal{A}_n(x)$ is freely generated as an R–module by the collection*

$$\{m_{\mathfrak{u}\mathfrak{v}} = v_\mathfrak{u}^* m_\lambda v_\mathfrak{v} : \textit{for } \mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\lambda) \textit{ and } \lambda \textit{ a partition of } n\}. \tag{2.1}$$

Moreover, the following statements hold.

1. *The R–linear map defined by $* : m_{\mathfrak{u}\mathfrak{v}} \mapsto m_{\mathfrak{v}\mathfrak{u}}$, for $\mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\lambda)$ and λ a partition of n, is the algebra anti–involution of $\mathcal{A}_n(x)$ satisfying $e_i \mapsto e_i$ for $i = 1, \ldots, n-1$.*
2. *Suppose that $b \in \mathcal{A}_n(x)$. If λ is a partition of n, and $\mathfrak{u} \in \mathfrak{T}_n(\lambda)$, then there exist $a_\mathfrak{v} \in R$, for $\mathfrak{v} \in \mathfrak{T}_n(\lambda)$, such that*

$$m_{\mathfrak{s}\mathfrak{u}} b \equiv \sum_{\mathfrak{v}} a_\mathfrak{v} m_{\mathfrak{s}\mathfrak{v}} \mod \check{\mathcal{A}}_n^\lambda, \tag{2.2}$$

for all $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$.

Note that Lemma 2.3 implies that, if λ is a partition of n, then $\check{\mathcal{A}}_n^\lambda$ is the R–module freely generated by the set $\{m_{\mathfrak{u}\mathfrak{v}} : \mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\mu), \text{ for } \mu \rhd \lambda\}$.

For k an integer with $1 \le 2k+1 < n$, and $\mu = (k, n-2k)$, let

$$\mathfrak{T}_n^{(k)}(\mu) = \{\mathfrak{s} \in \mathfrak{T}_n(\mu) : v_{\mathfrak{s}} \in \langle e_{2k+1}, \dots, e_{n-1}\rangle\}.$$

After observing that the map

$$* : m_{\mathfrak{u}\mathfrak{v}} \mapsto m_{\mathfrak{v}\mathfrak{u}}, \qquad \text{for } \mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\lambda), \text{ and } \lambda \text{ a partition of } n,$$

coincides with the algebra anti–involution that fixes the set $\{e_i : i = 1, \dots, n-1\}$ pointwise, Lemma 2.3 will follow from the following statement.

Lemma 2.4. *The set* $\{m_{\mathfrak{u}\mathfrak{v}} : \mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\lambda)$ *and* λ *a partition of* $n\}$ *freely generates* $\mathcal{A}_n(x)$ *as an* R*–module. Moreover, if* $b \in \mathcal{A}_n(x)$, $\lambda = (f, n-2f)$ *is a partition, and* $\mathfrak{u} \in \mathfrak{T}_n(\lambda)$, *then there exist* $a_{\mathfrak{v}} \in R$, *for* $\mathfrak{v} \in \mathfrak{T}_n(\lambda)$, *which depend only on* $\mathfrak{u}$, *such that*

$$m_\lambda v_{\mathfrak{u}} b = \sum_{\mathfrak{v} \in \mathfrak{T}_n(\lambda)} a_{\mathfrak{v}} m_\lambda v_{\mathfrak{v}} + \sum_{\substack{\mu \rhd \lambda \\ \mathfrak{r}, \mathfrak{t} \in \mathfrak{T}_n(\mu)}} a_{\mathfrak{r}\mathfrak{t}} m_{\mathfrak{r}\mathfrak{t}}, \tag{2.3}$$

where the sum is over partitions $\mu = (k, n-2k)$, *and* $\mathfrak{r} \in \mathfrak{T}_n^{(f)}(\mu)$, *with* $k = f+1, f+2, \dots,$ *and* $a_{\mathfrak{r}\mathfrak{t}} \in R$, *for* $\mathfrak{r} \in \mathfrak{T}_n^{(f)}(\mu)$ *and* $\mathfrak{t} \in \mathfrak{T}_n(\mu)$.

Lemma 2.5. *Let* $\lambda = (f, n-2f)$, *where* $n > n-2f > 0$. *Write* $\tau = (f+1, n-2f-2)$ *and* $\nu = (f, n-2f-1)$. *If* $\mathfrak{t} \in \mathfrak{T}_{n-1}^{(f-1)}(\nu)$ *and* $\mathfrak{u} \in \mathfrak{T}_{n-1}(\nu)$, *then either*

$$e_{2f-1} w_{2f,n} v_{\mathfrak{t}}^* m_\nu v_{\mathfrak{u}} = m_\lambda v_{\mathfrak{v}}, \qquad \textit{where } \mathfrak{v} \in \mathfrak{T}_n(\lambda) \textit{ and } \mathfrak{v}|_{n-1} = \mathfrak{u},$$

or there exists $\mathfrak{s} \in \mathfrak{T}_n^{(f)}(\tau)$, *such that*

$$e_{2f-1} w_{2f,n} v_{\mathfrak{t}}^* m_\nu v_{\mathfrak{u}} = v_{\mathfrak{s}}^* m_\tau v_{\mathfrak{v}}, \qquad \textit{where } \mathfrak{v} \in \mathfrak{T}_n(\tau) \textit{ and } \mathfrak{v}|_{n-1} = \mathfrak{u}.$$

Proof. We may write $v_{\mathfrak{t}}^* = w_{j,2f}$, where $2f \le j \le n-1$, so that

$$\begin{aligned} e_{2f-1} w_{2f,n} v_{\mathfrak{t}}^* m_\nu &= e_{2f-1} w_{2f,n} w_{j,2f} m_\nu \\ &= \begin{cases} e_{2f-1} e_{2f} m_\nu, & \text{if } j = n-1; \\ e_{2f-1} e_{2f} e_{j+1} e_{j+2} \cdots e_{n-1} m_\nu, & \text{if } 2f \le j < n-1. \end{cases} \end{aligned}$$

In the first case in the above expression, we obtain

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} = e_{2f-1} e_{2f} e_1 \cdots e_{2f-1} v_{\mathfrak{u}} = e_1 \cdots e_{2f-1} v_{\mathfrak{u}} = m_\lambda v_{\mathfrak{v}},$$

whereas in the second,

$$\begin{aligned} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} &= e_{2f-1} e_{2f} e_{j+1} e_{j+2} \cdots e_{n-1} (e_1 \cdots e_{2f-1}) v_{\mathfrak{u}} \\ &= e_1 \cdots e_{2f-1} e_{j+1} e_{j+2} \cdots e_{n-1} v_{\mathfrak{u}} \\ &= e_{j+1} e_j \cdots e_{2f+2} m_\tau e_{2f+2} e_{2f+3} \cdots e_{n-1} v_{\mathfrak{u}} = v_{\mathfrak{s}}^* m_\tau v_{\mathfrak{v}}, \end{aligned}$$

as required. □

If λ is a partition of n, let $m_{\mathfrak{t}^\lambda} = m_\lambda + \check{\mathcal{A}}_n^\lambda \in \mathcal{A}_n^\lambda / \check{\mathcal{A}}_n^\lambda$, and define C^λ to be the right $\mathcal{A}_n(x)$–submodule of $\mathcal{A}_n / \check{\mathcal{A}}_n^\lambda$ generated by $m_{\mathfrak{t}^\lambda}$. Further, if $\lambda = (f, n-2f)$ and $\mu \to \lambda$, define

$$y_\mu^\lambda = \begin{cases} m_{\mathfrak{t}^\lambda}, & \text{if } \mu = (f-1, n-2f+1); \\ m_{\mathfrak{t}^\lambda} w_{2f,n}, & \text{if } \mu = (f, n-2f-1), \end{cases}$$

and, let N^μ denote the $\mathcal{A}_{n-1}(x)$–submodule of C^λ generated by y_μ^λ.

In the next two lemmas, we assume that Lemma 2.4 is valid when applied to the algebra $\mathcal{A}_{n-1}(x)$ and show that the lemma is also true when applied to the algebra $\mathcal{A}_n(x)$ in the case that λ is maximal among partitions of n.

Lemma 2.6. *Let $n = 2f$ and $\lambda = (f, 0)$. If $\mu = (f-1, 1)$, then $\{y_\mu^\lambda v_\mathfrak{s} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$ generates C^λ as an R–module, and the R–module map $C^\lambda \to C^\mu$ determined by*

$$y_\mu^\lambda v_\mathfrak{s} \mapsto m_\mathfrak{t}, \qquad\qquad \text{for } \mathfrak{t} \in \mathfrak{T}_n(\lambda) \text{ and } \mathfrak{s} = \mathfrak{t}|_{n-1} \in \mathfrak{T}_{n-1}(\mu),$$

is an isomorphism of $\mathcal{A}_{n-1}(x)$–modules.

Proof. Let $b \in \mathcal{A}_{n-1}(x)$. Since $\check{\mathcal{A}}_{n-1}^\mu = 0$, by Lemma 2.4, which we apply inductively, there exist $a_\mathfrak{v} \in R$, for $\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)$, depending only on b, such that

$$m_\mu b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)} a_\mathfrak{v} m_\mu v_\mathfrak{v}.$$

Since $m_\lambda = e_{2f-1} m_\mu$, we multiply both sides of the above expression by e_{2f-1} on the left to obtain

$$m_\lambda b = e_{2f-1} m_\mu b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)} a_\mathfrak{v} e_{2f-1} m_\mu v_\mathfrak{v} = \sum_{\mathfrak{u} \in \mathfrak{T}_n(\lambda)} a_\mathfrak{u} m_\lambda v_\mathfrak{u},$$

where $a_\mathfrak{v} = a_\mathfrak{u}$ whenever $\mathfrak{u}|_{n-1} = \mathfrak{v}$.

Now we show that the collection $\{y_\mu^\lambda v_\mathfrak{s} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$ generates C^λ as an R–module. If $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, and $\mathfrak{s} = \mathfrak{t}|_{n-1}$, then either

(i) Shape$(\mathfrak{s}|_{n-2}) = (f-1, 0)$, in which event $v_\mathfrak{s} \in \mathcal{A}_{n-2}(x)$, or
(ii) Shape$(\mathfrak{s}|_{n-2}) = (f-2, 2)$, in which event $v_\mathfrak{t} = e_{n-2} v_\mathfrak{u}$, where $v_\mathfrak{u} \in \mathcal{A}_{n-2}(x)$.

In the case (i), $m_\lambda v_\mathfrak{s} e_{n-1} = x m_\lambda v_\mathfrak{s}$, while, in the case (ii), $m_\lambda v_\mathfrak{t} e_{n-1} = m_\lambda e_{n-2} v_\mathfrak{u} e_{n-1} = m_\lambda v_\mathfrak{u}$ which, since $v_\mathfrak{u} \in \mathcal{A}_{n-1}(x)$, can be written as

$$m_\lambda v_\mathfrak{u} = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)} a_\mathfrak{v} m_\lambda v_\mathfrak{v},$$

where $a_\mathfrak{v} \in R$, for $\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)$. Thus, if $b \in \mathcal{A}_{n-1}(x) e_{n-1} \mathcal{A}_{n-1}(x)$, then $m_\lambda b$ can be expressed as an R–linear combination of terms from $\{y_\mu^\lambda v_\mathfrak{s} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$. This completes the proof of the lemma. □

Lemma 2.7. *Let $n = 2f + 1$ and $\lambda = (f, 1)$. If $\mu^{(1)} = (f, 0)$ and $\mu^{(2)} = (f-1, 2)$, then*

$$(0) = N^{\mu^{(0)}} \subseteq N^{\mu^{(1)}} \subseteq N^{\mu^{(2)}} = C^\lambda \tag{2.4}$$

is a filtration of the $\mathcal{A}_n(x)$*–module* C^λ *by* $\mathcal{A}_{n-1}(x)$*–modules. Moreover, if* $\mu \in \{\mu^{(1)}, \mu^{(2)}\}$*, then* $\{y^\lambda_\mu v_\mathfrak{s} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$ *freely generates* N^μ *as an* R*–module, and the* R*–module homomorphism* $N^{\mu^{(i)}}/N^{\mu^{(i-1)}} \mapsto C^\lambda$ *determined by*

$$y^\lambda_\mu v_\mathfrak{s} + N^{\mu^{(i-1)}} \mapsto m_\mathfrak{t}, \qquad \text{for } \mathfrak{t} \in \mathfrak{T}_n(\lambda) \text{ and } \mathfrak{s} = \mathfrak{t}|_{n-1} \in \mathfrak{T}_{n-1}(\mu), \tag{2.5}$$

is an isomorphism of $\mathcal{A}_{n-1}(x)$*–modules.*

Proof. We have $y^\lambda_{\mu^{(1)}} = m_{\mathfrak{t}^\lambda}$ and $y^\lambda_{\mu^{(2)}} = m_{\mathfrak{t}^\lambda} e_{2f}$, so $y^\lambda_{\mu^{(2)}} e_{2f-1} = y^\lambda_{\mu^{(1)}}$, which shows that $N^{\mu^{(1)}} \subseteq N^{\mu^{(2)}}$ is an inclusion of $\mathcal{A}_{n-1}(x)$–modules. Furthermore, if $b \in \mathcal{A}_{n-1}(x)$ and $\mathfrak{u} \in \mathfrak{T}_{n-1}(\mu^{(1)})$, then, by Lemma 2.6, there exist $a_\mathfrak{v} \in R$, for $\mathfrak{v} \in \mathfrak{T}_{n-1}(\lambda)$, such that

$$m_{\mu^{(1)}} v_\mathfrak{u} b = \sum_{\mathfrak{v} \in \mathfrak{T}_n(\mu^{(1)})} a_\mathfrak{v} m_{\mu^{(1)}} v_\mathfrak{v};$$

thus,

$$y^\lambda_{\mu^{(1)}} v_\mathfrak{u} b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu^{(1)})} a_\mathfrak{v} y^\lambda_{\mu^{(1)}} v_\mathfrak{v}.$$

If $\mathfrak{u} \in \mathfrak{T}_{n-1}(\mu^{(2)})$, then $y^\lambda_{\mu^{(2)}} v_\mathfrak{u} b = e_{2f-1} e_{2f} m_{\mu^{(2)}} v_\mathfrak{u} b + \check{\mathcal{A}}^\lambda_{n-1}$, and, by Lemma 2.4 which we apply inductively,

$$e_{2f-1} e_{2f} m_{\mu^{(2)}} v_\mathfrak{u} b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu^{(2)})} a_\mathfrak{v} e_{2f-1} e_{2f} m_{\mu^{(2)}} v_\mathfrak{v} + \sum_{\mathfrak{r},\mathfrak{t} \in \mathfrak{T}_{n-1}(\mu^{(1)})} a_{\mathfrak{r}\mathfrak{t}} e_{2f-1} e_{2f} v^*_\mathfrak{r} m_{\mu^{(1)}} v_\mathfrak{t},$$

which shows that

$$y^\lambda_{\mu^{(2)}} v_\mathfrak{u} b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu^{(2)})} a_\mathfrak{v} y^\lambda_{\mu^{(2)}} v_\mathfrak{v} + \sum_{\mathfrak{r},\mathfrak{t} \in \mathfrak{T}_{n-1}(\mu^{(1)})} a_{\mathfrak{r}\mathfrak{t}} e_{2f-1} e_{2f} v^*_\mathfrak{r} m_{\mu^{(1)}} v_\mathfrak{t},$$

where the sum is over $\mathfrak{r} \in \mathfrak{T}^{(f-1)}_{n-1}(\mu^{(1)})$. Since in fact $v_\mathfrak{r} = 1$ whenever $\mathfrak{r} \in \mathfrak{T}^{(f-1)}_{n-1}(\mu^{(1)})$, and $e_{2f-1} e_{2f} m_{\mu^{(1)}} = m_\lambda = y^\lambda_{\mu^{(1)}}$, from the above expression, we obtain

$$y^\lambda_{\mu^{(2)}} v_\mathfrak{u} b = \sum_{\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu^{(2)})} a_\mathfrak{v} y^\lambda_{\mu^{(2)}} v_\mathfrak{v} + \sum_{\mathfrak{r},\mathfrak{t} \in \mathfrak{T}_{n-1}(\mu^{(1)})} a_\mathfrak{t} m_{\mu^{(1)}} v_\mathfrak{t},$$

which shows that the R–module map map $N^{\mu^{(2)}}/N^{\mu^{(1)}} \to C^{\mu^{(1)}}$ given by (2.5) is a homomorphism of $\mathcal{A}_{n-1}(x)$–modules. It remains to show that if $b \in \mathcal{A}_{n-1}(x) e_{n-1} \mathcal{A}_{n-1}(x)$, and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, then $m_\lambda v_\mathfrak{t} b$ can be expressed as an R–linear combination of elements from $\{m_\lambda v_\mathfrak{s} : \mathfrak{s} \in \mathfrak{T}_n(\lambda)\}$.

Let $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, $\mathfrak{t}|_{n-1} \in \mathfrak{T}_{n-1}(\mu^{(1)})$ and $\mathfrak{u} = \mathfrak{t}|_{n-2}$. Then $\text{Shape}(\mathfrak{u}) = (f-1, 1)$ and $v_\mathfrak{t} = v_\mathfrak{u} \in \mathcal{A}_{n-2}(x)$. Hence $v_\mathfrak{t} e_{2f} = e_{2f} v_\mathfrak{u} = v_\mathfrak{v}$, where $\mathfrak{v} \in \mathfrak{T}_n(\lambda)$ satisfies $\mathfrak{v}|_{n-1} \in \mathfrak{T}_{n-1}(\mu^{(2)})$; It follows that $m_\lambda v_\mathfrak{t} e_{n-1} = m_\lambda e_{2f} v_\mathfrak{t} = m_\lambda v_\mathfrak{v}$, which shows that in this instance, if $b \in \mathcal{A}_{n-1}(x) e_{n-1} \mathcal{A}_{n-1}(x)$, then $m_\lambda v_\mathfrak{t} b$ can be expressed as an R–linear combination of the required form.

If $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, $\mathfrak{t}|_{n-1} \in \mathfrak{T}_{n-1}(\mu^{(2)})$, and $\mathfrak{u} = \mathfrak{t}|_{n-2}$, then either

(i) $\text{Shape}(\mathfrak{u}) = (f-1, 1)$, in which event $v_\mathfrak{u} \in \mathcal{A}_{n-2}(x)$ and $v_\mathfrak{t} = e_{2f} v_\mathfrak{u}$, or

(ii) $\text{Shape}(\mathfrak{u}) = (f-2, 3)$, in which event $v_\mathfrak{t} = e_{2f} e_{2f-2} e_{2f-1} v_\mathfrak{u}$, where $v_\mathfrak{u} \in \mathcal{A}_{n-2}(x)$.

In the case (i) above,

$$m_\lambda v_\mathfrak{t} e_{n-1} = m_\lambda e_{2f} v_\mathfrak{u} e_{n-1} = m_\lambda e_{2f} v_\mathfrak{u} e_{2f} = x m_\lambda e_{2f} v_\mathfrak{u} = x m_\lambda v_\mathfrak{t},$$

while, in case (ii),

$$m_\lambda v_{\mathfrak{t}} e_{n-1} = m_\lambda e_{2f} e_{2f-2} e_{2f-1} v_{\mathfrak{u}} e_{n-1} = m_\lambda e_{2f} e_{2f-2} e_{2f-1} v_{\mathfrak{u}} e_{2f} = m_\lambda e_{2f} e_{2f-2} v_{\mathfrak{u}}.$$

Since $e_{2f-2} v_{\mathfrak{u}} \in \mathcal{A}_{n-1}(x)$, the rightmost term in the above equalities can be expressed as a linear combination of elements from $\{m_\lambda v_{\mathfrak{s}} : \mathfrak{s} \in \mathfrak{T}_n(\lambda)\}$. This completes the proof of the lemma. □

The following corollary provides the base case in the induction used in the proof of Lemma 2.9.

Corollary 2.8. *Let f be a non–negative integer, and $n = 2f + \delta$, where $\delta \in \{0, 1\}$ and $\lambda = (f, \delta)$. If Lemma 2.4 holds for $\mathcal{A}_{n-1}(x)$, then the set $\{m_{\mathfrak{u}\mathfrak{v}} = v_{\mathfrak{u}} m_\lambda v_{\mathfrak{v}} : \mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_k(\lambda)\}$ freely generates $\mathcal{A}_n^\lambda$ as an R–module. Furthermore, if $u \in \mathfrak{T}_n(\lambda)$ and $b \in \mathcal{A}_n(x)$, then there exist $a_{\mathfrak{v}} \in R$, for $\mathfrak{v} \in \mathfrak{T}_{n-1}(\lambda)$, depending only on $\mathfrak{u}$ and b, such that*

$$m_\lambda v_{\mathfrak{u}} b = \sum_{\mathfrak{v} \in \mathfrak{T}_n(\lambda)} a_{\mathfrak{v}} m_\lambda v_{\mathfrak{v}}.$$

In the next lemma, we take $\lambda = (f, n - 2f)$ to be a partition with $n - 2f > 1$, and, using Corollary 2.8, assume that Lemma 2.4 holds for for $\mathcal{A}_{n-1}(x)$ and for $\mathcal{A}_n(x)$ in the case of partitions $\nu \rhd \lambda$.

Lemma 2.9. *Let $\lambda = (f, n - 2f)$, where $n > n - 2f > 1$. If $\mu^{(1)} = (f, n - 2f - 1)$ and $\mu^{(2)} = (f - 1, n - 2f + 1)$. Then*

$$(0) = N^{\mu^{(0)}} \subseteq N^{\mu^{(1)}} \subseteq N^{\mu^{(2)}} = C^\lambda \tag{2.6}$$

is a filtration of the $\mathcal{A}_n(x)$–module C^λ by $\mathcal{A}_{n-1}(x)$–modules. Moreover, if $\mu \in \{\mu^{(1)}, \mu^{(2)}\}$, then $\{y_\mu^\lambda v_{\mathfrak{s}} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$ freely generates N^μ as an R–module, and the R–module homomorphism $N^{\mu^{(i)}}/N^{\mu^{(i-1)}} \mapsto C^\lambda$ determined by

$$y_\mu^\lambda v_{\mathfrak{s}} + N^{\mu^{(i-1)}} \mapsto m_{\mathfrak{t}}, \qquad \text{for } \mathfrak{t} \in \mathfrak{T}_n(\lambda) \text{ and } \mathfrak{s} = \mathfrak{t}|_{n-1} \in \mathfrak{T}_{n-1}(\mu), \tag{2.7}$$

is an isomorphism of $\mathcal{A}_{n-1}(x)$–modules.

Proof. First, if $\nu = (k, n - 2k - 1) \rhd \mu^{(1)}$, and $b \in \mathcal{A}_{n-1}^\nu$, then $C^\lambda b = 0$, since $\mathcal{A}_{n-1}^\nu \subset \check{\mathcal{A}}_n^\lambda$.

Next, observe that, if we write $\mu = (2f, n - 2f - 1)$, then, consistent with the inclusion of algebras in (1.4), the $\mathcal{A}_{n-1}(x)$–module N^μ is isomorphic to the $\mathcal{A}_{n-1}(x)$–module C^μ. Thus, by induction, $\{y_\mu^\lambda v_{\mathfrak{s}} : \mathfrak{s} \in \mathfrak{T}_{n-1}(\mu)\}$ freely generates N^μ as an R–module.

If $\mu = (f - 1, n - 2f + 1)$, then $y_\mu^\lambda w_{n-1,2f-1} = m_{\mathfrak{t}^\lambda}$, showing that (2.6) is an inclusion of $\mathcal{A}_{n-1}(x)$–modules.

Now, let $\mu = (f - 1, n - 2f - 1)$, suppose that $\mathfrak{t} \in \mathfrak{T}_{n-1}(\mu)$, and consider the action of an element $b \in \mathcal{A}_{n-1}(x)$ in the expression

$$m_\lambda w_{f,n} v_{\mathfrak{t}} b = e_{2f-1} w_{2f,n} m_\mu v_{\mathfrak{t}} b. \tag{2.8}$$

By Lemma 2.4, which we apply inductively, there exist $a_{\mathfrak{v}} \in R$, for $\mathfrak{v} \in \mathfrak{T}_n(\lambda)$, which depend only on $\mathfrak{t}$, such that

$$m_\mu v_{\mathfrak{t}} b = \sum_{\mathfrak{v} \in \mathfrak{T}_n(\mu)} a_{\mathfrak{v}} m_\mu v_{\mathfrak{v}} + \sum_{\substack{\nu \rhd \mu \\ \mathfrak{s}, \mathfrak{u} \in \mathfrak{T}_{n-1}(\nu)}} a_{\mathfrak{s}\mathfrak{u}} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}}, \tag{2.9}$$

where the latter sum is over partitions $\nu = (k, n-2k-1)$, for $k = f, f+1, \dots,$ and $\mathfrak{s} \in \mathfrak{T}_{n-1}^{(f-1)}(\nu)$. Substituted into (2.8), the expression (2.9) gives

$$\begin{aligned} m_\lambda w_{2f,n} v_{\mathfrak{t}} b &= \sum_{\mathfrak{v}\in\mathfrak{T}_n(\mu)} a_{\mathfrak{v}} e_{2f-1} w_{2f,n} m_\mu v_{\mathfrak{v}} + \sum_{\substack{\nu\rhd\mu\\ \mathfrak{s},\mathfrak{u}\in\mathfrak{T}_n(\nu)}} a_{\mathfrak{s}\mathfrak{u}} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} \\ &= \sum_{\mathfrak{v}\in\mathfrak{T}_n(\mu)} a_{\mathfrak{v}} m_\lambda w_{2f,n} v_{\mathfrak{v}} + \sum_{\substack{\nu\rhd\mu\\ \mathfrak{s},\mathfrak{u}\in\mathfrak{T}_n(\nu)}} a_{\mathfrak{s}\mathfrak{u}} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} \\ &= \sum_{\substack{\mathfrak{r}\in\mathfrak{T}_n(\lambda),\\ \mathfrak{r}|_{n-1}\in\mathfrak{T}_{n-1}(\mu)}} a_{\mathfrak{r}} m_\lambda v_{\mathfrak{r}} + \sum_{\substack{\nu\rhd\mu\\ \mathfrak{s},\mathfrak{u}\in\mathfrak{T}_n(\nu)}} a_{\mathfrak{s}\mathfrak{u}} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}}, \end{aligned} \tag{2.10}$$

where, in the above expression, $a_{\mathfrak{r}} \in R$ are defined, for $\mathfrak{r} \in \mathfrak{T}_n(\lambda)$, by the condition that $a_{\mathfrak{v}} = a_{\mathfrak{r}}$ whenever $\mathfrak{r}|_{n-1} = \mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)$.

Now, using Lemma 2.5, we turn our consideration to the summands $e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}}$ appearing in (2.10). Let $\tau = (f+1, n-2f-2)$; if $\nu = (f, n-2f-1)$, then either

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} = m_\lambda v_{\mathfrak{v}'}, \qquad \text{where } \mathfrak{v}' \in \mathfrak{T}_{n-1}(\lambda) \text{ and } \operatorname{Shape}(\mathfrak{v}'|_{n-1}) = \mu^{(1)}, \tag{2.11}$$

or, there exists $\mathfrak{t}' \in \mathfrak{T}_n^{(f)}(\tau)$, such that

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu v_{\mathfrak{u}} = v_{\mathfrak{t}'}^* m_\tau v_{\mathfrak{v}'}, \qquad \text{where } \mathfrak{v}' \in \mathfrak{T}_n(\tau) \text{ and } \mathfrak{v}'|_{n-1} = \mathfrak{u}. \tag{2.12}$$

Now suppose that $\nu = (k, n-2k-1)$, where $k = f+1, f+2, \dots$, let $\mathfrak{s} \in \mathfrak{T}_{n-1}^{(f-1)}(\nu)$, and consider the product $e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu$. We may write

$$v_{\mathfrak{s}}^* = w_{j_0,2i} w_{j_1,2i+2} \cdots w_{r,2k}, \qquad \text{where } f \le i \le k \text{ and } 2i \le j_0 < j_1 < \cdots \le r \le n-1.$$

If $v_{\mathfrak{s}}^* = 1$, then

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu = e_{2f-1} w_{2f,n} m_\nu = e_{2f-1} w_{2f,n} e_1 e_3 \cdots e_{2k-1} = w_{2k+2,n} e_1 e_3 \cdots e_{2k+1};$$

otherwise, if $f < i$, so that $v_{\mathfrak{s}}^* \in \langle e_{2f+2}, \dots, e_{n-2} \rangle$, then

$$\begin{aligned} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu &= e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* e_1 e_3 \cdots e_{2k-1} \\ &= e_{2f+1} w_{2f+2,n} v_{\mathfrak{s}}^* e_1 e_3 \cdots e_{2k-1} \\ &= e_{2f+1} w_{2f+2,n} v_{\mathfrak{s}}^* m_\nu. \end{aligned}$$

Thus we suppose that $v_{\mathfrak{s}}^* = w_{j_0,2f} w_{j_1,2f+2} \cdots w_{r,2k}$ where $2f < j_0 < j_1 < \cdots < r$, in which event,

$$w_{2f,n} w_{j_0,2f} = \begin{cases} e_{2f}, & \text{if } j_0 = n-1; \\ e_{2f} e_{j_0+1} e_{j_0+2} \cdots e_{n-1}, & \text{if } 2f < j_0 < n-1. \end{cases} \tag{2.13}$$

Let $v_{\mathfrak{s}'}^* = w_{j_1,2f+2} \cdots w_{r,2k}$; in the first case in (2.13), using $e_{2f-1} e_{2f} e_{2f-1} = e_{2f-1}$,

$$e_{2f} w_{2f,n} v_{\mathfrak{s}}^* m_\nu = e_{2f-1} w_{2f,n} w_{n-1,2f} v_{\mathfrak{s}'}^* m_\nu = e_{2f-1} e_{2f} v_{\mathfrak{s}'}^* m_\nu = v_{\mathfrak{s}'}^* m_\nu,$$

and in the second,

$$\begin{aligned} e_{2f-1} w_{2f,n} v_{\mathfrak{s}}^* m_\nu &= e_{2f-1} e_{2f} e_{j_0+1} e_{j_0+2} \cdots e_{n-1} v_{\mathfrak{s}'}^* m_\nu \\ &= e_{j_0+1} e_{j_0+2} \cdots e_{n-1} v_{\mathfrak{s}'}^* m_\nu. \end{aligned}$$

Let $\nu' = (k, n-2k)$, so that $m_\nu \mapsto m_{\nu'}$ under the inclusion $\mathcal{A}_{n-1}(x) \hookrightarrow \mathcal{A}_n(x)$, and let $b' = e_{2f-1} w_{2f,n} v_{\mathfrak{s}}$ take a value in $\{v_{\mathfrak{s}'}^*, e_{j_0+1} e_{j_0+2} \cdots e_{n-1} v_{\mathfrak{s}'}^*\}$, depending on the case in (2.13). Since $b' \in \langle e_i : 2f < i < n \rangle$, and $\nu' \rhd \lambda$, there exist $a_{\mathfrak{s}''} \in R$, for $\mathfrak{s}'' \in \mathfrak{T}_n(\nu')$, such that

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}} m_\nu = b' m_{\nu'} = \sum_{\mathfrak{s}'' \in \mathfrak{T}_n^{(f)}(\nu')} v_{\mathfrak{s}''}^* m_{\nu'} + \sum_{\substack{\tau' \rhd \nu' \\ \mathfrak{r}', \mathfrak{t}' \in \mathfrak{T}_n(\tau')}} a_{\mathfrak{r}'\mathfrak{t}'} v_{\mathfrak{r}'}^* m_{\tau'} v_{\mathfrak{t}'}, \tag{2.14}$$

where $\mathfrak{r}' \in \mathfrak{T}_n^{(f)}(\tau')$, $\mathfrak{t}' \in \mathfrak{T}_n^{(k)}(\tau')$, and $a_{\mathfrak{r}'\mathfrak{t}'} \in R$, for $\tau' \rhd \nu'$. If $\mathfrak{u} \in \mathfrak{T}_{n-1}(\nu)$, then $v_{\mathfrak{u}} = v_{\mathfrak{u}'}$, where $\mathfrak{u}'|_{n-1} = \mathfrak{u} \in \mathfrak{T}_{n-1}(\nu)$ and $\mathfrak{u} \in \mathfrak{T}_n(\nu')$, we multiply both sides of (2.14) by $v_{\mathfrak{u}} = v_{\mathfrak{u}'}$ to obtain

$$e_{2f-1} w_{2f,n} v_{\mathfrak{s}} m_\nu v_{\mathfrak{u}} = \sum_{\mathfrak{s}'' \in \mathfrak{T}_n^{(f)}(\nu')} v_{\mathfrak{s}''}^* m_{\nu'} v_{\mathfrak{u}'} + \sum_{\substack{\tau' \rhd \nu' \\ \mathfrak{r}'', \mathfrak{t}'' \in \mathfrak{T}_n(\tau')}} a_{\mathfrak{r}''\mathfrak{t}''} v_{\mathfrak{r}''}^* m_{\tau'} v_{\mathfrak{t}''}, \tag{2.15}$$

where $\mathfrak{r}'' \in \mathfrak{T}_n^{(f)}(\tau')$, $\mathfrak{t}'' \in \mathfrak{T}_n(\tau')$, and $a_{\mathfrak{r}''\mathfrak{t}''} \in R$, for $\tau' \rhd \nu'$.

Combining (2.10) with (2.11), (2.12) and (2.15), we have shown that if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, $\operatorname{Shape}(\mathfrak{t}|_{n-1}) = \mu^{(2)}$ and $b \in \mathcal{A}_{n-1}(x)$, then there exist $a_{\mathfrak{r}}, a_{\mathfrak{v}'} \in R$, for $\mathfrak{r}, \mathfrak{v}' \in \mathfrak{T}_n(\lambda)$, where $\operatorname{Shape}(\mathfrak{r}|_{n-1}) = \mu^{(2)}$ and $\operatorname{Shape}(\mathfrak{v}'|_{n-1}) = \mu^{(1)}$, satisfying

$$m_\lambda v_{\mathfrak{t}} b = \sum_{\substack{\mathfrak{r} \in \mathfrak{T}_n(\lambda) \\ \operatorname{Shape}(\mathfrak{r}|_{n-1}) = \mu^{(2)}}} a_{\mathfrak{r}} m_\lambda v_{\mathfrak{r}} + \sum_{\substack{\mathfrak{v}' \in \mathfrak{T}_n(\lambda) \\ \operatorname{Shape}(\mathfrak{v}'|_{n-1}) = \mu^{(1)}}} a_{\mathfrak{v}'} m_\lambda v_{\mathfrak{v}'} + \sum_{\substack{\gamma \rhd \lambda \\ \mathfrak{s}, \mathfrak{u} \in \mathfrak{T}_n(\gamma)}} a_{\mathfrak{s}\mathfrak{u}} v_{\mathfrak{s}}^* m_\gamma v_{\mathfrak{u}}, \tag{2.16}$$

where the sum is over $\mathfrak{s} \in \mathfrak{T}_n^{(f)}(\gamma)$, $\mathfrak{u} \in \mathfrak{T}_n(\gamma)$, and $a_{\mathfrak{s}\mathfrak{u}} \in R$, for $\gamma \rhd \lambda$. The manner in which the $a_{\mathfrak{r}} \in R$, for $\mathfrak{r} \in \mathfrak{T}_n(\lambda)$ satisfying $\operatorname{Shape}(\mathfrak{r}|_{n-1}) = \mu^{(2)}$, are derived in (2.10) from the action of $\mathcal{A}_{n-1}(x)$ on $C^{\mu^{(2)}}$ shows that the map (2.7) is a homomorphism of $\mathcal{A}_{n-1}(x)$–modules.

It remains to demonstrate that $N^{\mu^{(2)}} = C^\lambda$. To this purpose, we show that if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ and $b \in \mathcal{A}_{n-1}(x) e_{n-1} \mathcal{A}_{n-1}(x)$, then $m_\lambda v_{\mathfrak{t}} b$ can be expressed as a sum of the form (2.16). Firstly, we suppose that $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ and let $\mathfrak{u} = \mathfrak{t}_{n-1}$ satisfy $\operatorname{Shape}(\mathfrak{u}) = \mu^{(1)}$. In this case, $v_{\mathfrak{t}} = v_{\mathfrak{u}} \in \mathcal{A}_{n-1}(x)$. If $v_{\mathfrak{u}} \in \mathcal{A}_{n-2}(x)$, then

$$m_\lambda v_{\mathfrak{t}} e_{n-1} = m_\lambda e_{n-1} v_{\mathfrak{t}} = w_{n,2f+2} m_\nu w_{2f+2,n} v_{\mathfrak{t}}, \quad \text{where } \nu = (f+1, n-2f-2). \tag{2.17}$$

By what we have already shown, the term appearing on the right hand side of (2.17) can be written as a sum of the form (2.16). Otherwise, if $v_{\mathfrak{u}} = w_{2f,n-1} v_{\mathfrak{v}}$, where $\operatorname{Shape}(\mathfrak{v}) = (f-1, n-2f)$ and $v_{\mathfrak{v}} \in \mathcal{A}_{n-2}(x)$, then

$$m_\lambda v_{\mathfrak{t}} e_{n-1} = m_\lambda w_{2f,n-1} v_{\mathfrak{v}} e_{n-1} = m_\lambda w_{2f,n-1} e_{n-1} v_{\mathfrak{v}} = m_\lambda w_{2f,n} v_{\mathfrak{v}} = m_\lambda v_{\mathfrak{s}} \tag{2.18}$$

where $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$ is defined by $\mathfrak{s}|_{n-2} = \mathfrak{v}$ and $\operatorname{Shape}(\mathfrak{s}|_{n-1}) = (f-1, n-2f+1)$. Now suppose that $v_{\mathfrak{t}} = w_{2f,n} v_{\mathfrak{u}}$, where $\operatorname{Shape}(\mathfrak{u}) = (f-1, n-2f+1)$. If $v_{\mathfrak{u}} \in \mathcal{A}_{n-2}(x)$, then $m_\lambda v_{\mathfrak{t}} e_{n-1} = x m_\lambda v_{\mathfrak{t}}$; otherwise $v_{\mathfrak{u}} = w_{2f-2,n-1} v_{\mathfrak{v}}$, where $\operatorname{Shape}(\mathfrak{v}) = (f-2, n-2f+2)$ and $v_{\mathfrak{v}} \in \mathcal{A}_{n-2}(x)$. Thus,

$$m_\lambda v_{\mathfrak{t}} e_{n-1} = m_\lambda w_{2f,n} w_{2f-2,n-1} v_{\mathfrak{v}} e_{n-1} = m_\lambda w_{2f,n} w_{2f-2,n} v_{\mathfrak{v}} = m_\lambda w_{2f,n} w_{2f-2,n-2} v_{\mathfrak{v}},$$

which is a term that we have already shown can be expressed as a sum of the form (2.16). This completes the proof of the lemma. □

Proof of Lemma 2.4. Firstly, if $b \in \mathcal{A}_{n-1}(x)$, then (2.3) holds by virtue of the calculations preceding (2.16) in the proof of Lemma 2.8 and, if $b \in \mathcal{A}_{n-1}(x) e_{n-1} \mathcal{A}_{n-1}(x)$, then the proof of the fact that $N^{\mu^{(2)}} = C^\lambda$ in the proof of Lemma 2.8 shows that (2.3) holds. □

Example 2.1. If $n = 6$ and $\lambda = (2,2)$, then the elements $m_\lambda v_\mathfrak{s}$, for $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$, are given in terms of the diagram presentation for $\mathcal{A}_n(x)$ are as follows:

$\mathfrak{s} = ((0,0),(0,1),(1,0),(1,1),(2,0),(2,1),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(1,1),(2,0),(2,1),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(1,0),(1,1),(1,2),(2,1),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(1,1),(1,2),(2,1),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(0,3),(1,2),(2,1),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(1,0),(1,1),(1,2),(1,3),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(1,1),(1,2),(1,3),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(0,3),(1,2),(1,3),(2,2))$: $m_\lambda v_\mathfrak{s} =$



$\mathfrak{s} = ((0,0),(0,1),(0,2),(0,3),(0,4),(1,3),(2,2))$: $m_\lambda v_\mathfrak{s} =$



3. Jucys–Murphy Elements for Temperley–Lieb Algebras

In [4], T. Halverson, M. Mazzocco and A. Ram have defined a family of commuting operators in the affine Temperley–Lieb algebras. The operators of [4] are analogues to the Jucys–Murphy elements from the representation theory of the symmetric group. For the purposes of these notes, it will be useful to define in $\mathcal{A}_n(x)$ a sequence $(T_i : i = 0, 1, \dots)$ by $T_0 = 0$, $T_1 = 0$, $T_2 = e_1$, and

$$T_{i+1} = -e_iT_i - T_ie_i + e_ie_{i-1}T_ie_i - z_{i-1} - e_iz_{i-2}, \qquad \text{for } i = 2, 3, \dots$$

where, $z_1 = 0$, and

$$z_i = \sum_{k=0}^{i} x^k T_{i-k}, \qquad \text{for } i = 2, 3, \dots.$$

Lemma 3.1. *For $i = 2, 3, \dots$, the following statements hold:*

(1) $e_{i+1}e_iT_ie_{i+1} = e_{i+1}T_ie_ie_{i+1}$;

(2) $e_ie_{i-1}T_ie_i = e_iT_ie_{i-1}e_i$;
(3) $z_i^* = z_i$ *and* $T_i^* = T_i$;
(4) $e_{i-1}T_{i+1} = T_{i+1}e_{i-1}$;
(5) $e_i(xT_i + T_{i+1}) = (xT_i + T_{i+1})e_i$;
(6) T_i *commutes with* $\mathcal{A}_{i-1}(x)$;
(7) *the element* $z_i = \sum_{k=0}^{i} x^kT_{i-k}$ *is central in* $\mathcal{A}_i(x)$.

Proof. The item (1) follows from the relation $e_{i+1}e_ie_{i+1} = e_{i+1}$ and fact that e_{i+1} commutes with T_i. Turning to the statement (2), which is true when $i = 2$, we proceed by induction. Since

$$T_i = -e_{i-1}T_{i-1} - T_{i-1}e_{i-1} + e_{i-1}e_{i-2}T_{i-1}e_{i-1} - z_{i-2} - z_{i-3}e_{i-1},$$

applying (1) yields

$$\begin{aligned} e_ie_{i-1}T_ie_i &= -xe_ie_{i-1}T_{i-1}e_i - e_ie_{i-1}T_{i-1}e_{i-1}e_i \\ &\qquad + xe_ie_{i-1}e_{i-2}T_{i-1}e_{i-1}e_i - e_ie_{i-1}z_{i-2}e_i - xe_ie_{i-1}z_{i-3}e_i \\ &= xe_iT_{i-1} - e_ie_{i-1}T_{i-1}e_{i-1}e_i \\ &\qquad + xe_ie_{i-1}T_{i-1}e_{i-2}e_{i-1}e_i - e_ie_{i-1}e_iz_{i-2} - xe_ie_{i-1}e_iz_{i-3} \\ &= xe_iT_{i-1} - e_ie_{i-1}T_{i-1}e_{i-1}e_i \\ &\qquad + xe_ie_{i-1}e_{i-2}T_{i-1}e_{i-1}e_i - e_iz_{i-2} - xe_iz_{i-3} \\ &= e_iT_ie_{i-1}e_i, \end{aligned}$$

as required. The statement (3) follows from (2), while

$$\begin{aligned} e_{i-1}T_{i+1} &= -e_{i-1}e_iT_i - e_{i-1}T_ie_i + e_{i-1}e_ie_{i-1}T_ie_i - e_{i-1}z_{i-1} - e_{i-1}z_{i-2}e_i \\ &= -e_{i-1}e_iT_i - e_{i-1}T_ie_i + e_{i-1}T_ie_i - e_{i-1}z_{i-1} - e_{i-1}z_{i-2}e_i \\ &= -e_{i-1}e_iT_i - e_{i-1}T_{i-1} - e_{i-1}z_{i-1} - e_{i-1}z_{i-2}e_i. \end{aligned} \tag{3.1}$$

Now,

$$\begin{aligned} e_{i-1}e_iT_i + e_{i-1}T_{i-1} &= -e_{i-1}e_ie_{i-1}T_{i-1} - e_{i-1}e_iT_{i-1}e_{i-1} \\ &\qquad + e_{i-1}e_ie_{i-1}e_{i-2}T_{i-1}e_{i-1} - e_{i-1}e_iz_{i-2} - e_{i-1}e_iz_{i-3}e_{i-1} + e_{i-1}T_{i-1} \\ &= -e_{i-1}T_{i-1} - e_{i-1}e_iT_{i-1}e_{i-1} \\ &\qquad + e_{i-1}e_{i-2}T_{i-1}e_{i-1} - e_{i-1}e_iz_{i-2} - e_{i-1}z_{i-3} + e_{i-1}T_{i-1} \\ &= -e_{i-1}e_iT_{i-1}e_{i-1} + e_{i-1}e_{i-2}T_{i-1}e_{i-1} - e_{i-1}e_iz_{i-2} - e_{i-1}z_{i-3}, \end{aligned}$$

which, substituted into (3.1), yields

$$\begin{aligned} e_{i-1}T_{i+1} &= e_{i-1}e_iT_{i-1}e_{i-1} - e_{i-1}e_{i-2}T_{i-1}e_{i-1} + e_{i-1}z_{i-2}e_i - e_{i-1}z_{i-2}e_i + e_{i-1}z_{i-3} \\ &= e_{i-1}e_iT_{i-1}e_{i-1} - e_{i-1}e_{i-2}T_{i-1}e_{i-1} + e_{i-1}z_{i-3} \\ &= e_{i-1}T_{i-1}e_ie_{i-1} - e_{i-1}T_{i-1}e_{i-2}e_{i-1} + e_{i-1}z_{i-3} \\ &= T_{i+1}e_{i-1}. \end{aligned}$$

To see (5), we have

$$\begin{aligned}e_i(xT_i+T_{i+1}) &= xe_iT_i - xe_iT_i - e_iT_ie_i + xe_ie_{i-1}T_ie_i - e_iz_{i-1} - xz_{i-2}e_i\\ &= -e_iT_ie_i + xe_ie_{i-1}T_ie_i - e_iz_{i-1} - xz_{i-2}e_i\\ &= -e_iT_ie_i + xe_iT_ie_{i-1}e_i - z_{i-1}e_i - xz_{i-2}e_i\\ &= (xT_i+T_{i+1})e_i.\end{aligned}$$

The proof of (6) and (7) is a joint induction. We assume that e_k commutes with T_i whenever $k = 1, \dots, i-2$, and that z_k is central in $\mathcal{A}_k(x)$ whenever $k = 1, 2, \dots, i$. Since it is already known that T_{i+1} commutes with e_{i-1}, we first show that T_{i+1} commutes with e_{i-2}. By item (4),

$$\begin{aligned}T_{i+1}e_{i-2} &= -e_iT_ie_{i-2} - T_ie_ie_{i-2} + e_iT_ie_{i-1}e_ie_{i-2} - z_{i-1}e_{i-2} - e_iz_{i-2}e_{i-2}\\ &= -e_{i-2}e_iT_i - e_{i-2}T_ie_i + e_iT_ie_{i-1}e_ie_{i-2} - e_{i-2}z_{i-1} - e_iz_{i-2}e_{i-2}.\end{aligned} \tag{3.2}$$

Hence we must show that $e_iT_ie_{i-1}e_i - e_iz_{i-2}$ commutes with e_{i-2}; to keep the indices within nice bounds, we demonstrate that $e_{k+1}T_{k+1}e_ke_{k+1} - e_{k+1}z_{k-1}$ commutes with e_{k-1}:

$$\begin{aligned}e_{k+1}T_{k+1}e_ke_{k+1}e_{k-1} - e_{k+1}z_{k-1}e_{k-1} &= -e_{k+1}e_kT_ke_ke_{k+1}e_{k-1} - xe_{k+1}T_ke_ke_{k+1}e_{k-1}\\ &\quad + xe_{k+1}e_kT_ke_{k-1}e_ke_{k+1}e_{k-1} - e_{k+1}z_{k-1}e_ke_{k+1}e_{k-1}\\ &\quad - xe_{k+1}e_kz_{k-2}e_{k+1}e_{k-1} - e_{k+1}z_{k-1}e_{k-1}\\ &= -e_{k+1}e_kT_ke_ke_{k+1}e_{k-1} - xe_{k+1}T_ke_{k-1} + xe_{k+1}T_ke_{k-1} - e_{k+1}z_{k-1}e_{k-1}\\ &\quad - xe_{k+1}z_{k-2}e_{k-1} - e_{k+1}z_{k-1}e_{k-1}\\ &= -e_{k+1}e_kT_ke_ke_{k+1}e_{k-1} - xe_{k+1}e_{k-1}z_{k-2} - 2e_{k+1}z_{k-1}e_{k-1}.\end{aligned}$$

Since $(e_{k+1}e_{k-1}z_{k-2})^* = e_{k+1}e_{k-1}z_{k-2}$, we consider

$$\begin{aligned}e_{k+1}e_kT_ke_ke_{k+1}e_{k-1} + 2e_{k+1}z_{k-1}e_{k-1} &= -e_{k+1}e_ke_{k-1}T_{k-1}e_ke_{k+1}e_{k-1}\\ &\quad - e_{k+1}e_kT_{k-1}e_{k-1}e_ke_{k+1}e_{k-1} + e_{k+1}e_ke_{k-1}T_{k-1}e_{k-2}e_{k-1}e_ke_{k+1}e_{k-1} - e_{k+1}e_kz_{k-2}e_{k+1}e_{k-1}\\ &\quad - e_{k+1}e_ke_{k-1}z_{k-3}e_ke_{k+1}e_{k-1} + 2e_{k+1}z_{k-1}e_{k-1}\\ &= -e_{k+1}T_{k-1}e_{k-1} - e_{k+1}T_{k-1}e_{k-1} + e_{k+1}e_{k-1}T_{k-1}e_{k-2}e_{k-1} - xe_{k+1}z_{k-2}e_{k-1} - e_{k+1}e_{k-1}z_{k-3}\\ &\quad + 2e_{k+1}z_{k-1}e_{k-1}\\ &= e_{k+1}e_{k-1}T_{k-1}e_{k-2}e_{k-1} - xe_{k+1}z_{k-2}e_{k-1} - 2e_{k+1}z_{k-2}e_{k-1}\\ &= e_{k+1}e_{k-1}e_{k-2}T_{k-1}e_{k-1} - xe_{k-1}e_{k+1}z_{k-2} - 2e_{k-1}e_{k+1}z_{k-2},\end{aligned}$$

which shows that $e_{k+1}e_kT_ke_ke_{k+1}e_{k-1} + 2e_{k+1}z_{k-1}e_{k-1}$ is fixed by $* : \mathcal{A}_{k+2}(z) \to \mathcal{A}_{k+2}(z)$, and therefore that $e_iT_ie_{i-1}e_i - e_iz_{i-2}$ commutes with e_{i-2}. Hence, by (3.2), T_{i+1} commutes with e_{i-2}. Now, if $k = 1, \dots, i-3$, then by induction,

$$\begin{aligned}T_{i+1}e_k &= -e_iT_ie_k - T_ie_ie_k + e_ie_{i-1}T_ie_ie_k - z_ie_k - e_iz_{i-2}e_k\\ &= -e_ke_iT_i - e_kT_ie_i + e_ke_ie_{i-1}T_ie_i - e_kz_i - e_ke_iz_{i-2} = e_kT_{i+1}.\end{aligned}$$

The item (5) shows that z_{i+1} commutes with $e_1, \dots, e_i$, which completes the induction and the proof of the lemma. □

Since the elements of $(T_i : i = 0, 1, \dots)$ commute, we are justified in referring to the T_i as "Jucys–Murphy elements" for the Temperley–Lieb algebras.

Define a sequence $(p_i \in R : i = 0, 1, \dots)$ by $p_0 = 0$, $p_1 = 1$ and $p_{i+1} = xp_i - p_{i-1}$ for $i = 1, 2, \dots$. For $k = 1, 2, \dots$, introduce a sequence $(T_i^{(k)} : i = 0, 1, \dots)$ by $T_0^{(k)} = 0$, $T_1^{(k)} = 0$, $T_2^{(k)} = e_k$, and, for $i = 2, 3, \dots$,

$$T_{i+1}^{(k)} = -e_{k+i-1}T_i^{(k)} - T_i^{(k)}e_{k+i-1} + e_{k+i-1}e_{k+i-2}T_i^{(k)}e_{k+i-1} - z_{i-1}^{(k)} - z_{i-2}^{(k)}e_{k+i-1},$$

where $z_1^{(k)} = 0$, and

$$z_i^{(k)} = \sum_{j=0}^{i} x^j T_{i-j}^{(k)}, \qquad \text{for } i = 2, 3, \dots,$$

so that the $T_i^{(k)}$, for $i = 0, 1, \dots$, and $k = 1$, are just the Jucys–Murphy elements.

Lemma 3.2. *For $i = 1, 2, \dots$, $z_i e_1 = e_1 z_{i-2}^{(3)} + p_i e_1$, and $T_{i+1}e_1 = e_1 T_{i-1}^{(3)} - p_{i-1}e_1$.*

Proof. Note that $e_1 z_2 = p_2 e_1 + e_1 z_1^{(3)}$ and $e_1 T_3 = e_1 T_1^{(3)} - p_1 e_1$, and proceed by induction. If $i \geq 3$, then

$$\begin{aligned}
T_{i+1}e_1 &= -e_i T_i e_1 - T_i e_1 e_i + e_i e_{i-1} T_i e_1 e_i - z_{i-1}e_1 - e_i z_{i-2} e_1 \\
&= -e_1 e_i T_{i-2}^{(3)} - e_1 T_{i-2}^{(3)} e_i + 2p_{i-2}e_1 e_i + e_1 e_i e_{i-1} T_{i-2}^{(3)} e_i - p_{i-2} e_1 e_i e_{i-1} e_i \\
&\quad - e_1 z_{i-3}^{(3)} - p_{i-1}e_1 - e_1 e_i z_{i-4}^{(3)} - p_{i-2} e_1 e_i \\
&= -e_1 e_i T_{i-2}^{(3)} - e_1 T_{i-2}^{(3)} e_i + e_1 e_i e_{i-1} T_{i-2}^{(3)} e_i - e_1 z_{i-3}^{(3)} - e_1 e_i z_{i-4}^{(3)} - p_{i-1} e_1 \\
&= e_1 T_{i-1}^{(3)} - p_{i-1} e_1,
\end{aligned}$$

while

$$\begin{aligned}
z_i e_1 = x z_{i-1} e_1 + T_i e_1 &= x e_1 z_{i-3}^{(3)} + x p_i e_1 + e_1 T_{i-1}^{(3)} - p_{i-1} e_1 \\
&= e_1 (x z_{i-3}^{(3)} + T_{i-2}^{(3)}) + (x p_{i-1} - p_{i-2}) e_1 = e_1 z_{i-2}^{(3)} + p_i e_1,
\end{aligned}$$

as required. □

Corollary 3.3. *For $i = 1, 2, \dots$, and $k = 1, 2, \dots$,*

$$z_i^{(k)} e_k = e_k z_{i-2}^{(k+2)} + p_i e_k, \qquad \text{and} \qquad T_{i+1}^{(k)} e_k = e_k T_{i-1}^{(k+2)} - p_{i-1} e_k.$$

The following elementary lemma will be used to give the eigenvalues of the Jucys–Murphy elements.

Lemma 3.4. *The sequence $(p_i \in R : i = 0, 1, \dots)$ satisfies the following relations:*

$$p_{2j} + p_{2j+2} + \cdots + p_{2i} = p_{i-k+1}p_{i+k} - p_{j-k}p_{j+k+1}, \qquad \text{for } k \in \{1, 2, \dots, j\}; \tag{3.3}$$

$$p_{2j+1} + p_{2j+3} + \cdots + p_{2i+1} = p_{i-k+1}p_{i+k+1} - p_{j-k}p_{j+k}, \qquad \text{for } k \in \{0, 1, \dots, j\}. \tag{3.4}$$

Proof. We first show that,

$$p_{2i} = p_{i-k+1}p_{i+k} - p_{i-k}p_{i+k-1}, \qquad \text{for } k = 1, \dots, i. \tag{3.5}$$

$$p_{2i+1} = p_{i-k+1}p_{i+k+1} - p_{i-k}p_{i+k}, \qquad \text{for } k = 0, \dots, i. \tag{3.6}$$

Since (3.5) and (3.6) both hold when $k = i$, by induction,

$$\begin{aligned} p_{i-k+2}p_{i+k-1} - p_{i-k+1}p_{i+k-2} &= (xp_{i-k+1} - p_{i-k})p_{i+k-1} - p_{i-k+1}p_{i+k-2} \\ &= p_{i-k+1}(xp_{i+k-1} - p_{i-k-2}) - p_{i-k}p_{i+k-1} \\ &= p_{i-k+1}p_{i+k} - p_{i-k}p_{i+k-1} = p_{2i}, \end{aligned}$$

and,

$$\begin{aligned} p_{i-k+2}p_{i+k} - p_{i-k+1}p_{i+k-1} &= (xp_{i-k+1} - p_{i-k})p_{i+k} - p_{i-k+1}p_{i+k-1} \\ &= p_{i-k+1}(xp_{i+k} - p_{i-k-1}) - p_{i-k}p_{i+k} \\ &= p_{i-k+1}p_{i+k+1} - p_{i-k}p_{i+k} = p_{2i+1}. \end{aligned}$$

From (3.5), we have, for $k \in \{1, \dots, j\}$, a telescoping sum

$$\begin{aligned} p_{2j} + p_{2j+2} + \cdots + p_{2i} = p_{j-k+1}p_{j+k} - p_{j-k}p_{j+k-1} \\ + p_{j-k+2}p_{j+k+1} - p_{j-k+1}p_{j+k} + \cdots + p_{i-k+1}p_{i+k} - p_{i-k}p_{i+k-1} \\ = p_{i-k+1}p_{i+k} - p_{j-k}p_{j+k-1}. \end{aligned}$$

Similarly, from (3.6), we have, for $k \in \{0, \dots, j\}$, a sum

$$\begin{aligned} p_{2j+1} + p_{2j+3} + \cdots + p_{2i+1} = p_{j-k+1}p_{j+k+1} - p_{j-k}p_{j+k} + \\ + p_{j-k+2}p_{j+k+2} - p_{j-k+1}p_{j+k+1} + \cdots + p_{i-k+1}p_{i+k+1} - p_{i-k}p_{i+k} \\ = p_{i-k+1}p_{i+k+1} - p_{j-k}p_{j+k}. \end{aligned}$$

□

It will be useful to note that from the previous lemma:

$$\begin{aligned} &p_4 + p_6 + \cdots + p_{2i} = p_{i-1}p_{i+2} = (xp_i - p_{i+1})p_{i+2}; \\ &p_{2j} + p_{2j+2} + \cdots + p_{2i} = p_{i-j+1}p_{i+j} && \text{for } j = 1, 2, \dots, i-1; \\ &p_{2j+1} + p_{2j+3} + \cdots + p_{2i+1} = p_{i-j+1}p_{i+j+1} && \text{for } j = 0, 1, \dots, i-1. \end{aligned}$$

Lemma 3.5. *If $\lambda = (f, n-2f)$ is a partion of n, and i is an integer, $1 \le i \le n$, then*

$$m_{\mathfrak{t}^\lambda} z_i = \begin{cases} p_f p_{i-f+1} m_{\mathfrak{t}^\lambda}, & \text{if } 2f+1 \le i \le n; \\ p_k p_{i-k+1} m_{\mathfrak{t}^\lambda}, & \text{if } i = 2k, \text{ or } i = 2k+1, \text{ and } 0 \le k \le f. \end{cases}$$

Proof. The statement being true when $i \le 1$, we proceed by induction, first considering the case where $i \in \{2k, 2k+1 : 0 \le k \le f\}$. If $2 = k \le f$ and and z_{2k-1} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar $p_{k-1}p_{k+1}$, then

$$m_\lambda T_{2k} = e_1 e_3 T_{2k} = e_1 e_3 T_2^{(3)} - p_2 e_1 e_3 = 0,$$

so that, in this instance, $z_{2k} = (xz_{2k-1} + T_{2k})$ acts by $xp_{k-1}p_{k+1} = p_k p_{k+1}$. If $2 < k \le f$, and z_{2k-1} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar $p_{k-1}p_{k+1}$, then

$$\begin{aligned} e_1 e_3 \cdots e_{2k-1} T_{2k} &= e_1 e_3 \cdots e_{2k-1} T_4^{(2k-3)} - (p_4 + p_6 + \cdots + p_{2k-2}) e_1 e_3 \cdots e_{2k-1} \\ &= -(p_4 + p_6 + \cdots + p_{2k-2}) e_1 e_3 \cdots e_{2k-1} \\ &= (p_k p_{k+1} - xp_{k-1}p_{k+1}) e_1 e_3 \cdots e_{2k-1}, \end{aligned}$$

so that

$$m_{\mathfrak{t}^\lambda} z_{2k} = m_{\mathfrak{t}^\lambda}(xz_{2k-1} + T_{2k}) = (xp_{k-1}p_{k+1} + p_k p_{k+1} - xp_{k-1}p_{k+1})m_{\mathfrak{t}^\lambda} = p_k p_{k+1} m_{\mathfrak{t}^\lambda}.$$

If $0 \leq k \leq f$, and z_{2k} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar $p_k p_{k+1}$, then

$$\begin{aligned} e_1 e_3 \cdots e_{2k-1} T_{2k+1} &= e_1 e_3 \cdots e_{2k-1} T_1^{(2k+1)} - (p_1 + p_3 + \cdots + p_{2k-1}) e_1 e_3 \cdots e_{2k-1} \\ &= (p_k p_{k+2} - xp_k p_{k+1}) e_1 e_3 \cdots e_{2k-1}, \end{aligned}$$

so that

$$m_{\mathfrak{t}^\lambda} z_{2k+1} = m_{\mathfrak{t}^\lambda}(xz_{2k} + T_{2k+1}) = (xp_k p_{k+1} + p_k p_{k+2} - xp_k p_{k+1}) m_{\mathfrak{t}^\lambda} = p_k p_{k+2} m_{\mathfrak{t}^\lambda}.$$

Now we turn our attention to the action of z_i case where $2f+1 \leq i \leq n$. If $i = 2k$, where $f < k$, and z_{i-1} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar $p_f p_{2k-f}$, then

$$\begin{aligned} m_\lambda T_{2k} = e_1 e_3 \cdots e_{2f-1} T_{2k} &= e_1 e_3 \cdots e_{2f-1} T_{2k-2f}^{(2f+1)} - \sum_{j=k-f}^{k-1} p_{2j} e_1 e_3 \cdots e_{2f-1} \\ &\equiv -\sum_{j=k-f}^{k-1} p_{2j} m_\lambda = -p_f p_{2k-f-1} m_\lambda \mod \check{\mathcal{A}}_n^\lambda, \end{aligned}$$

so z_{2k} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar

$$xp_f p_{2k-f} - p_f p_{2k-f-1} = p_f(xp_{2k-f} - p_{2k-f-1}) = p_f p_{2k-f+1} = p_f p_{i-f+1}.$$

Similarly, if $i = 2k+1$, and z_{i-1} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar $p_f p_{2k-f+1}$ then

$$\begin{aligned} m_\lambda T_{2k+1} = e_1 e_3 \cdots e_{2f-1} T_{2k+1} &= e_1 e_3 \cdots e_{2f-1} T_{2k-2f+1}^{(2f+1)} - \sum_{j=k-f}^{k-1} p_{2j+1} e_1 e_3 \cdots e_{2f-1} \\ &\equiv -\sum_{j=k-f}^{k-1} p_{2j+1} m_\lambda = -p_f p_{2k-f} m_\lambda \mod \check{\mathcal{A}}_n^\lambda, \end{aligned}$$

so z_{2k+1} acts on $m_{\mathfrak{t}^\lambda}$ by the scalar

$$xp_f p_{2k-f+1} - p_f p_{2k-f} = p_f(xp_{2k-f+1} - p_{2k-f}) = p_f p_{2k-f+2} = p_f p_{i-f+1}.$$

□

Let λ be a partition of n and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$. Define a sequence $(r_\mathfrak{t}(k) \in R : k = 0, 1, \ldots, n)$ by $r_\mathfrak{t}(0) = 0$, $r_\mathfrak{t}(1) = 0$, and

$$r_\mathfrak{t}(k) = \begin{cases} (p_i - xp_{i-1})p_{k-i+1}, & \text{if } \operatorname{Shape}(\mathfrak{t}|_k) = (i, k-2i), \\ & \qquad \text{and } \operatorname{Shape}(\mathfrak{t}|_{k-1}) = (i-1, k-2i+1); \\ (p_{k-i+1} - xp_{k-i})p_i, & \text{if } \operatorname{Shape}(\mathfrak{t}|_k) = (i, k-2i), \\ & \qquad \text{and } \operatorname{Shape}(\mathfrak{t}|_{k-1}) = (i, k-2i-1), \end{cases}$$

for $k = 2, 3, \ldots, n$.

Corollary 3.6. *Let λ be a partition of n and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$. If k is an integer, $0 \leq k \leq n$, then there exist $a_\mathfrak{v} \in R$, for $\mathfrak{v} \in \mathfrak{T}_n(\lambda)$, such that*

$$m_\mathfrak{t} T_k = r_\mathfrak{t}(k) m_\mathfrak{t} + \sum_{\mathfrak{v} \rhd \mathfrak{t}} a_\mathfrak{v} m_\mathfrak{v}.$$

If λ is a partition of n and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, define

$$z_{\mathfrak{t}}(k) = \sum_{i=0}^{k} x^i r_{\mathfrak{t}}(k-i), \qquad \text{for } k = 0, 1, \dots, n.$$

Corollary 3.7. *Let λ be a partition of n and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$. If $\mu = (i, k-2i)$ and $\mathrm{Shape}(\mathfrak{t}|_k) = \mu$, then $z_{\mathfrak{t}}(k) = p_i p_{k-i+1}$, and there exist $a_{\mathfrak{s}} \in R$, for $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$, such that*

$$m_{\mathfrak{t}} z_k = z_{\mathfrak{t}}(k) m_{\mathfrak{t}} + \sum_{\mathfrak{s} \rhd \mathfrak{t}} a_{\mathfrak{s}} m_{\mathfrak{s}}.$$

Lemma 3.8. *Let λ be a partition of n and $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$. If $\mathfrak{s}|_{n-1} = \mathfrak{t}|_{n-1}$ and $r_{\mathfrak{s}}(n) = r_{\mathfrak{t}}(n)$, then $\mathfrak{s} = \mathfrak{t}$.*

4. An Orthogonal Basis

To determine the action of the generators of $\mathcal{A}_n(x)$ on an orthogonal basis for $\mathcal{A}_n(x)$, we require an alternative definition of the Jucys–Murphy elements as given in Corollary 4.2.

Lemma 4.1. *For $i = 2, 3, \dots,$*

$$e_i e_{i-1} T_i e_i = -x z_{i-1} e_i + z_{i-2} e_i + p_i e_i, \qquad \textit{and,} \qquad e_i T_i e_i = -2 z_{i-1} e_i + p_{i-1} e_i.$$

Proof. The lemma being true when $i = 2$, we proceed by induction:

$$\begin{aligned} e_{i+1} e_i T_{i+1} e_{i+1} &= -x e_{i+1} e_i T_i e_{i+1} - e_{i+1} e_i T_i e_i e_{i+1} + x e_{i+1} e_i e_{i-1} T_i e_i e_{i+1} - z_{i-1} e_{i+1} - x e_{i+1} z_{i-2} \\ &= -x T_i e_{i+1} + 2 z_{i-1} e_{i+1} - p_{i-1} e_{i+1} - x^2 z_{i-1} e_{i+1} + x z_{i-2} e_{i+1} + x p_i e_{i+1} - z_{i-1} e_{i+1} - x z_{i-2} e_{i+1} \\ &= -x z_i e_{i+1} + z_{i-1} e_{i+1} + p_{i+1} e_{i+1}, \end{aligned}$$

while

$$\begin{aligned} e_{i+1} T_{i+1} e_{i+1} &= -2 T_i e_{i+1} + e_{i+1} e_i e_{i-1} T_i e_i e_{i+1} - x z_{i-1} e_{i+1} - z_{i-2} e_{i+1} \\ &= -2 T_i e_{i+1} - x z_{i-1} e_{i+1} + z_{i-2} e_{i+1} + p_i e_{i+1} - x z_{i-1} e_{i+1} - z_{i-2} e_{i+1} \\ &= -2 z_i e_{i+1} + p_i e_{i+1}. \end{aligned}$$

□

Corollary 4.2. *For $i = 2, 3, \dots,$*

$$T_{i+1} = -e_i T_i - T_i e_i + (p_i - x z_{i-1}) e_i - z_{i-1}.$$

Corollary 4.3. *For $i = 2, 3, \dots,$ and $k = 1, 2, \dots,$*

$$e_i T_{i+1}^k = e_i (p_{i+1} - x z_i + z_{i-1})^k \tag{4.1}$$

Proof. From Corollary (4.2),

$$\begin{aligned} e_i T_{i+1} + x e_i T_i &= -e_i T_i e_i - x^2 z_{i-1} e_i - z_{i-1} e_i + x p_i e_i \\ &= 2 z_{i-1} e_i - p_{i-1} e_i - x^2 z_{i-1} e_i - z_{i-1} e_i + x p_i e_i, \end{aligned}$$

which shows that the statement (4.1) is true when $k = 1$. The general case now follows.

□

In what follows, we let F denote the field of fractions of R, and denote by $\mathscr{A}_n(x)$ the F–algebra generated by $e_1, \dots, e_n$, so that $\mathscr{A}_n(x) = \mathcal{A}_n(x) \otimes_R F$. Following [1], let

$$\mathfrak{R}(k) = \{r_{\mathfrak{t}}(k) : \mathfrak{t} \in \mathfrak{T}_k(\lambda) \text{ for } \lambda \text{ a partion of } n\}.$$

Let λ be a partition of n. As in [1], if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, define

$$F_{\mathfrak{t}} = \prod_{k=2}^{n} \prod_{\substack{r \in \mathfrak{R}(k), \\ r \neq r_{\mathfrak{s}}(k)}} \frac{T_k - r}{r_{\mathfrak{s}}(k) - r},$$

and let $f_{\mathfrak{t}} = m_{\mathfrak{t}} F_{\mathfrak{t}}$.

Lemma 4.4. *If λ is a partition of n, then*

1. *if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$, then there exist $a_{\mathfrak{u}} \in F$, for $\mathfrak{u} \in \mathfrak{T}_n(\lambda)$, such that*
$$f_{\mathfrak{t}} = m_{\mathfrak{t}} + \sum_{\mathfrak{u} \rhd \mathfrak{t}} a_{\mathfrak{u}} m_{\mathfrak{u}};$$
2. *the set $\{f_{\mathfrak{t}} : \mathfrak{t} \in \mathfrak{T}_n(\lambda)\}$ is a basis over F for the $\mathscr{A}_n(x)$–module C^{λ};*
3. *if $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$, then $\langle f_{\mathfrak{s}}, f_{\mathfrak{t}} \rangle = \delta_{\mathfrak{s},\mathfrak{t}}$;*
4. *if $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ and k is an integer, $0 \le k \le n$, then $f_{\mathfrak{t}} T_k = r_{\mathfrak{t}}(k) f_{\mathfrak{t}}$ and $f_{\mathfrak{t}} z_k = z_{\mathfrak{t}}(k) f_{\mathfrak{t}}$.*

Let λ be a partition of n. If $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ and k is an integer $1 \le k \le n$, define a set $\{e_k(\mathfrak{s}, \mathfrak{t}) \in F : \mathfrak{s} \in \mathfrak{T}_n(\lambda)\}$ by

$$f_{\mathfrak{t}} e_k = \sum_{\mathfrak{s} \in \mathfrak{T}_n(\lambda)} e_k(\mathfrak{s}, \mathfrak{t}) f_{\mathfrak{s}}. \tag{4.2}$$

Lemma 4.5. *Let λ be a partition of n and $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$. If $e_k(\mathfrak{s}, \mathfrak{t})$ are determined by* (4.2), *then*

1. *if $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$ and $e_k(\mathfrak{s}, \mathfrak{t}) \neq 0$, then $r_{\mathfrak{s}}(i) = r_{\mathfrak{t}}(i)$ whenever $i \neq k-1$ and $i \neq k$;*
2. *if $e_k(\mathfrak{s}, \mathfrak{t}) \neq 0$ for some $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$, then $\mathfrak{t}^{(k-1)} = \mathfrak{t}^{(k+1)}$ and $\mathfrak{s}^{(k-1)} = \mathfrak{s}^{(k+1)}$;*
3. *if $e_k(\mathfrak{s}, \mathfrak{t}) \neq 0$ for some $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$ with $\mathfrak{s} \neq \mathfrak{t}$, then $e_k(\mathfrak{u}, \mathfrak{t}) = 0$ whenever $\mathfrak{u} \notin \{\mathfrak{s}, \mathfrak{t}\}$;*
4. *if $e_k(\mathfrak{s}, \mathfrak{t}) \neq 0$, where $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$, and $\mathfrak{t} \rhd \mathfrak{s}$, then*

$$e_k(\mathfrak{t}, \mathfrak{t}) = \frac{r_{\mathfrak{t}}(k+1) + z_{\mathfrak{t}}(k-1)}{p_k - x z_{\mathfrak{t}}(k-1) - 2 r_{\mathfrak{t}}(k)}, \qquad \textit{and} \qquad e_k(\mathfrak{s}, \mathfrak{t}) = 1;$$
$$e_k(\mathfrak{s}, \mathfrak{s}) = \frac{r_{\mathfrak{s}}(k+1) + z_{\mathfrak{s}}(k-1)}{p_k - x z_{\mathfrak{s}}(k-1) - 2 r_{\mathfrak{s}}(k)}, \qquad \textit{and} \qquad e_k(\mathfrak{t}, \mathfrak{s}) = e_k(\mathfrak{s}, \mathfrak{s}) e_k(\mathfrak{t}, \mathfrak{t}).$$

Proof. The item (1) follows from the fact that e_k commutes with T_i whenever $i \notin \{k-1, k\}$. For the item (2), we use the fact (*cf.* Lemma 3.2) that

$$e_k z_{k+1} = e_k (z_{k-1} + p_{k+1})$$

to observe that

$$z_{\mathfrak{t}}(k+1) \sum_{\mathfrak{s} \in \mathfrak{T}_n(\lambda)} e_k(\mathfrak{s}, \mathfrak{t}) f_{\mathfrak{s}} = \sum_{\mathfrak{s} \in \mathfrak{T}_n(\lambda)} (z_{\mathfrak{s}}(k-1) + p_{k+1}) f_{\mathfrak{s}}. \tag{4.3}$$

If $\mathfrak{s} = (\lambda^{(0)}, \dots, \lambda^{(n)})$ and $\operatorname{Shape}(\mathfrak{s}|_{k-1}) = (i-1, k+1-2i)$, then there are four possibilities for the sequence $(\lambda^{(k-1)}, \lambda^{(k)}, \lambda^{(k+1)})$ given as follows:

$$(\lambda^{(k-1)}, \lambda^{(k)}, \lambda^{(k+1)}) = ((i-1, k-2i+1), (i, k-2i), (i, k-2i+1)) \tag{4.4}$$

$$(\lambda^{(k-1)}, \lambda^{(k)}, \lambda^{(k+1)}) = ((i-1, k-2i+1), (i-1, k-2i+2), (i, k-2i+1)) \tag{4.5}$$

$$(\lambda^{(k-1)}, \lambda^{(k)}, \lambda^{(k+1)}) = ((i-1, k-2i+1), (i, k-2i), (i+1, k-2i-1)) \tag{4.6}$$

$$(\lambda^{(k-1)}, \lambda^{(k)}, \lambda^{(k+1)}) = ((i-1, k-2i+1), (i, k-2i+2), (i, k-2i+3)). \tag{4.7}$$

Taking $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$ as in (4.6) and (4.7) respectively, the expression (4.3) gives:

$$e_k(\mathfrak{s}, \mathfrak{t}) p_{i+1} p_{k-i+1} = e_k(\mathfrak{s}, \mathfrak{t})(p_{i-1} p_{k-i+1} + p_{k+1}); \tag{4.8}$$

$$e_k(\mathfrak{s}, \mathfrak{t}) p_{i-1} p_{k-i+2} = e_k(\mathfrak{s}, \mathfrak{t})(p_{i-1} p_{k-i+1} + p_{k+1}); \tag{4.9}$$

Given that

$$p_{k+1} = p_{i+1} p_{k-i+2} - p_{i-1} p_{k-i+1}, \qquad \text{for } k = 1, 2, \dots, \text{ and } i = 1, \dots, k,$$

the statements (4.8) and (4.9) imply respectively that

$$p_{i+1}(p_{k-i+2} - p_{k-i+1}) e_k(\mathfrak{s}, \mathfrak{t}) = 0, \qquad \text{and} \qquad p_{k-i+1}(p_{i+1} - p_i) e_k(\mathfrak{s}, \mathfrak{t}) = 0,$$

conclusions which are patently absurd unless $e_k(\mathfrak{s}, \mathfrak{t}) = 0$ whenever $\mathfrak{s}^{(k-1)} \neq \mathfrak{s}^{(k+1)}$. The statement (3) now follows.

For the statement (4), we have

$$f_\mathfrak{t} T_{k+1} = -f_\mathfrak{t} e_k T_k - r_\mathfrak{t}(k) f_\mathfrak{t} e_k + (p_k - x z_\mathfrak{t}(k-1)) f_\mathfrak{t} e_k - z_\mathfrak{t}(k-1) f_\mathfrak{t}$$

which, comparing the coefficient of $f_\mathfrak{t}$ on both sides, implies that

$$r_\mathfrak{t}(k+1) = -r_\mathfrak{t}(k) e_k(\mathfrak{t}, \mathfrak{t}) - r_\mathfrak{t}(k) e_k(\mathfrak{t}, \mathfrak{t}) + (p_k - x z_\mathfrak{t}(k-1)) e_k(\mathfrak{t}, \mathfrak{t}) - z_\mathfrak{t}(k-1).$$

Now the fact that $r_\mathfrak{t}(k+1) + z_\mathfrak{t}(k-1) \neq 0$ shows that the stated expression for $e_k(\mathfrak{t}, \mathfrak{t})$ holds; the same reasoning yields the stated expression for $e_k(\mathfrak{s}, \mathfrak{s})$. To observe that $e_k(\mathfrak{s}, \mathfrak{t}) = 1$, by the maximality property of $\mathfrak{t}$ and Corollary 2.2,

$$f_\mathfrak{t} e_k = m_\mathfrak{t} e_k + \sum_{\mathfrak{u} \rhd \mathfrak{t}} a_\mathfrak{u} f_\mathfrak{u} e_k = m_\mathfrak{t} e_k = m_\mathfrak{s} = f_\mathfrak{s} - \sum_{\mathfrak{v} \rhd \mathfrak{s}} \alpha_\mathfrak{v} f_\mathfrak{v},$$

for some $a_\mathfrak{u}, \alpha_\mathfrak{v} \in F$, with $\mathfrak{u}, \mathfrak{v} \in \mathfrak{T}_n(\lambda)$.

To complete the proof of the lemma, $f_\mathfrak{t} e_k = e_k(\mathfrak{t}, \mathfrak{t}) f_\mathfrak{t} + f_\mathfrak{s}$ implies that

$$x f_\mathfrak{t} e_k = f_\mathfrak{t} e_k^2 = e_k(\mathfrak{t}, \mathfrak{t})(e_k(\mathfrak{t}, \mathfrak{t}) f_\mathfrak{t} + f_\mathfrak{s}) + e_k(\mathfrak{t}, \mathfrak{s}) f_\mathfrak{t} + e_k(\mathfrak{s}, \mathfrak{s}) f_\mathfrak{s},$$

whence, comparing coefficients,

$$x = e_k(\mathfrak{t}, \mathfrak{t}) + e_k(\mathfrak{s}, \mathfrak{s}), \qquad \text{and} \qquad x e_k(\mathfrak{t}, \mathfrak{t}) = (e_k(\mathfrak{t}, \mathfrak{t}))^2 + e_k(\mathfrak{t}, \mathfrak{s}).$$

Thus

$$e_k(\mathfrak{t}, \mathfrak{s}) = e_k(\mathfrak{t}, \mathfrak{t})(x - e_k(\mathfrak{t}, \mathfrak{t})) = e_k(\mathfrak{s}, \mathfrak{s}) e_k(\mathfrak{t}, \mathfrak{t}).$$

□

Corollary 4.6. *Let λ be a partition of n, and k be an integer, $1 \leq k < n$. Suppose that $\mathfrak{s}, \mathfrak{t} \in \mathfrak{T}_n(\lambda)$ satisfy $\mathfrak{t} \rhd \mathfrak{s}$ and $e_k(\mathfrak{s}, \mathfrak{t}) \neq 0$. If $\operatorname{Shape}(\mathfrak{t}|_k) = (i, k-2i)$, then*

$$e_k(\mathfrak{t}, \mathfrak{t}) = \frac{p_{k-2i+1}}{p_{k-2i+2}}, \qquad \textit{and} \qquad e_k(\mathfrak{s}, \mathfrak{s}) = \frac{p_{k-2i+3}}{p_{k-2i+2}}.$$

Proof. From (4.4) and (4.5), we have

$$r_{\mathfrak{t}}(k) = (p_i - xp_{i-1})p_{k-i+1}, \quad r_{\mathfrak{t}}(k+1) = (p_{k-i+2} - xp_{k-i+1})p_i, \quad z_{\mathfrak{t}}(k-1) = p_{i-1}p_{k-i+1}$$
$$r_{\mathfrak{s}}(k) = (p_{k-i+2} - xp_{k-i+1})p_{i-1}, \quad r_{\mathfrak{t}}(k+1) = (p_i - xp_{i-i})p_{k-i+2}, \qquad z_{\mathfrak{t}}(k-1) = p_{i-1}p_{k-i+1}.$$

Substituting the above into the expressions provided in item (4) of Lemma 4.5,

$$e_k(\mathfrak{t},\mathfrak{t}) = \frac{(p_{k-i+2} - xp_{k-i+1})p_i + p_{i-1}p_{k-i+1}}{p_k - xp_{i-1}p_{k-i+1} - 2(p_i - xp_{i-1})p_{k-i+1}} = \frac{p_ip_{k-i+2} - p_{i+1}p_{k-i+1}}{p_k + xp_{i-1}p_{k-i+1} - 2p_ip_{k-i+1}},$$

and

$$e_k(\mathfrak{s},\mathfrak{s}) = \frac{(p_i - xp_{i-i})p_{k-i+2} + p_{i-1}p_{k-i+1}}{p_k - xp_{i-1}p_{k-i+1} - 2(p_{k-i+2} - xp_{k-i+1})p_{i-1}} = \frac{p_ip_{k-i+2} - p_{i-1}p_{k-i+3}}{p_k + xp_{i-1}p_{k-i+1} - 2p_{i-1}p_{k-i+2}}.$$

The required formulae now follow from elementary considerations, namely, if $2 \le 2i \le k$,

$$p_{k-2i+1} = p_{i+1}p_{k-i+1} - p_ip_{k-i+2}, \quad \text{and} \quad p_{k-2i+2} = 2p_ip_{k-i+1} - xp_{i-1}p_{k-i+1} - p_k;$$
$$p_{k-2i+3} = p_ip_{k-i+2} - p_{i-1}p_{k-i+3}, \quad \text{and} \quad p_{k-2i+2} = p_k + xp_{i-1}p_{k-i+1} - 2p_{i-1}p_{k-i+2}.$$

□

Lemma 4.7. *Let λ be a partition of n, and k be an integer $1 \le k < n$. Suppose that $\mathfrak{s},\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ satisfy $\mathfrak{t} \rhd \mathfrak{s}$ and $e_k(\mathfrak{s},\mathfrak{t}) \neq 0$. If* $\operatorname{Shape}(\mathfrak{t}|_k) = (i, k-2i)$*, then*

$$\langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle = \frac{p_{k-2i+1}(xp_{k-2i+2} - p_{k-2i+1})}{(p_{k-2i+2})^2}\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle = \frac{p_{k-2i+1}p_{k-2i+3}}{(p_{k-2i+2})^2}\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle.$$

Proof. Since $f_{\mathfrak{t}}e_k = e_k(\mathfrak{t},\mathfrak{t})f_{\mathfrak{t}} + f_{\mathfrak{s}}$,

$$\langle f_{\mathfrak{t}}e_k, f_{\mathfrak{t}}e_k\rangle = (e_k(\mathfrak{t},\mathfrak{t}))^2\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle + \langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle,$$

while associativity of the bilinear form implies that

$$\langle f_{\mathfrak{t}}e_k, f_{\mathfrak{t}}e_k\rangle = x\langle f_{\mathfrak{t}}e_k, f_{\mathfrak{t}}\rangle = xe_k(\mathfrak{t},\mathfrak{t})\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle.$$

Thus $\langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle = e_k(\mathfrak{t},\mathfrak{t})(x - e_k(\mathfrak{t},\mathfrak{t}))\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle$, and the result follows from Corollary 4.6. □

5. THE DETERMINANT OF THE GRAM MATRIX

If λ is a partition of n, let $\dim(\lambda) = \sharp\{\mathfrak{t} : \mathfrak{t} \in \mathfrak{T}_n(\lambda)\}$ and write $\det(\lambda) = \prod_{\mathfrak{t}\in\mathfrak{T}_n(\lambda)}\langle f_{\mathfrak{t}}, f_{\mathfrak{t}}\rangle$ for the determinant of the Gram matrix associated with C^λ. The next result is a branching law.

Lemma 5.1. *Let $\lambda = (f, n-2f)$ and $\mu = (f-1, n-2f+1)$ be partitions, where $n - 2f \ge 0$ and $f \ge 1$. Then*

$$\det(\lambda) = \prod_{\nu\to\lambda}\det(\nu)\cdot\left(\frac{p_{n-2f+2}}{p_{n-2f+1}}\right)^{\dim(\mu)}$$

Proof. The result will follow once we show that if $\mathfrak{s} \in \mathfrak{T}_n(\lambda)$ and $\mathfrak{u} = \mathfrak{s}|_{n-1}$, then

$$\langle f_{\mathfrak{s}}, f_{\mathfrak{s}}\rangle = \langle f_{\mathfrak{u}}, f_{\mathfrak{u}}\rangle\frac{p_{n-2f+2}}{p_{n-2f+1}}, \qquad \text{whenever } \operatorname{Shape}(\mathfrak{u}) = \mu. \tag{5.1}$$

To prove the statement (5.1), we first consider the case where $\mathfrak{u} = \mathfrak{t}^\mu$. To this purpose define a sequence $(v_{\mathfrak{s}_i} \in \mathfrak{T}_n(\lambda) : i = 0, \dots, n-2f)$ by $v_{\mathfrak{s}_i} = w_{2f,2f+i}$, for $0 \le i \le n-2f$.

It follows that $\mathrm{Shape}(\mathfrak{s}_i|_{2f+i-1}) = (f-1, i+1)$ and $\mathrm{Shape}(\mathfrak{s}_i|_{2f+i}) = (f, i)$, while $\mathfrak{s}_i \rhd \mathfrak{s}_{i+1}$ and $e_{2f+i}(\mathfrak{s}_{i+1}, \mathfrak{s}_i) \neq 0$ for $0 \leq i < n-2f$. Hence

$$\langle f_{\mathfrak{s}_{i+1}}, f_{\mathfrak{s}_{i+1}} \rangle = \langle f_{\mathfrak{s}_i}, f_{\mathfrak{s}_i} \rangle \frac{p_{i+1} p_{i+3}}{(p_{i+2})^2},$$

and

$$\langle f_{\mathfrak{s}_{n-2f}}, f_{\mathfrak{s}_{n-2f}} \rangle = \langle f_{\mathfrak{s}_0}, f_{\mathfrak{s}_0} \rangle \frac{p_1 p_3}{(p_2)^2} \frac{p_2 p_4}{(p_3)^2} \frac{p_3 p_5}{(p_4)^2} \cdots \frac{p_{n-2f} p_{n-2f+2}}{(p_{n-2f+1})^2} = \langle f_{\mathfrak{s}_0}, f_{\mathfrak{s}_0} \rangle \frac{p_1}{p_2} \frac{p_{n-2f+2}}{p_{n-2f+1}}.$$

Since $\mathfrak{s} = \mathfrak{s}_{n-2f}$, and $\langle f_{\mathfrak{s}_0}, f_{\mathfrak{s}_0} \rangle = p_2 \langle f_{\mathfrak{u}}, f_{\mathfrak{u}} \rangle$, the above verifies (5.1) when $\mathfrak{u} = \mathfrak{t}^\mu$.

Now suppose that $\mathfrak{s}|_{n-1} = \mathfrak{u} \in \mathfrak{T}_{n-1}(\mu)$, and that $\mathfrak{t}^\mu \rhd \mathfrak{u}$. Then there exists $\mathfrak{v} \in \mathfrak{T}_{n-1}(\mu)$ such that $\mathfrak{v} \rhd \mathfrak{u}$ and $v_{\mathfrak{u}} = v_{\mathfrak{v}} e_k$, for some k with $1 \leq k < n$. If $\mathfrak{t} \in \mathfrak{T}_n(\lambda)$ satisfies $\mathfrak{t}|_{n-1} = \mathfrak{v}$, then $\mathfrak{t} \rhd \mathfrak{s}$ and

$$\langle f_{\mathfrak{s}}, f_{\mathfrak{s}} \rangle = \langle f_{\mathfrak{t}}, f_{\mathfrak{t}} \rangle \frac{p_{k-2i+1} p_{k-2i+3}}{(p_{k-2i+2})^2}, \qquad \text{where } (i, k-2i) = \mathrm{Shape}(\mathfrak{t}|_k)$$

while, by induction,

$$\langle f_{\mathfrak{t}}, f_{\mathfrak{t}} \rangle = \langle f_{\mathfrak{v}}, f_{\mathfrak{v}} \rangle \frac{p_{n-2f+2}}{p_{n-2f+1}}.$$

Since

$$\langle f_{\mathfrak{u}}, f_{\mathfrak{u}} \rangle = \langle f_{\mathfrak{v}}, f_{\mathfrak{v}} \rangle \frac{p_{k-2i+1} p_{k-2i+3}}{(p_{k-2i+2})^2}, \qquad \text{where } (i, k-2i) = \mathrm{Shape}(\mathfrak{v}|_k),$$

it follows that (5.1) holds in general. □

If $\lambda = (i, n-2i)$, and $\mu = (j, n-2j)$ are partitions, with $\mu \rhd \lambda$, define

$$g_{\lambda,\mu} = \left[\frac{p_{n-i-j+1}}{p_{i-j}} \right]^{\dim(\mu)}$$

Since the dimensions of modules C^μ are given in terms of certain binomial coefficients [5], the next statement gives closed formulae for the Gram determinants associated with the Temperley–Lieb algebras (*cf.* Corollary 4.7 of [3]).

Lemma 5.2. *Let $\lambda = (f, n-2f)$ be a partition. If $n-2f > 0$, then*

$$\det(\lambda) = \prod_{\mu \rhd \lambda} g_{\lambda,\mu},$$

and, if $n-2f = 0$, then $\det(\lambda) = \det(\mu) \cdot x^{\dim(\lambda)}$, where $\mu = (f-1, n-2f+1)$.

Proof. We first assume that $n-2f > 0$. Let

$$\lambda^{(i)} = (f-i, n-2f+2i), \quad \text{and} \quad \mu^{(i)} = (f-i, n-2f+2i-1), \quad \text{for } i = 0, 1, \ldots, f.$$

Then $\lambda = \lambda^{(0)} \rhd \lambda^{(1)} \rhd \cdots \rhd \lambda^{(f)}$ and $\mu = \mu^{(0)} \rhd \mu^{(1)} \rhd \cdots \rhd \mu^{(f)}$, while $\mu^{(i)} \to \lambda^{(i)}$ for $i = 0, \ldots, f$, and $\mu^{(i+1)} \to \lambda^{(i)}$ for $i = 0, \ldots, f-1$. If $n-2f > 1$, then from Lemma 5.1 and induction,

$$\begin{aligned} \det(\lambda) &= \det\big(\mu^{(0)}\big) \det\big(\mu^{(1)}\big) \Big(\frac{p_{n-2f+2}}{p_{n-2f+1}} \Big)^{\dim(\mu^{(1)})} \\ &= \prod_{i=1}^{f} g_{\mu^{(0)},\mu^{(i)}} \cdot \prod_{j=1}^{f-1} g_{\mu^{(1)},\mu^{(j+1)}} \cdot \Big(\frac{p_{n-2f+2}}{p_{n-2f+1}} \Big)^{\dim(\mu^{(1)})}, \end{aligned}$$

where, for $i = 1, \dots, f$, and $j = 1, \dots, f-1$,

$$g_{\mu^{(0)},\mu^{(i)}} = \Big(\frac{p_{n-2f+i}}{p_i}\Big)^{\dim(\mu^{(i)})} \qquad \text{and} \qquad g_{\mu^{(1)},\mu^{(j+1)}} = \Big(\frac{p_{n-2f+j+2}}{p_j}\Big)^{\dim(\mu^{(j+1)})}.$$

Thus,

$$\begin{aligned}
\det(\lambda) &= \prod_{i=1}^{f}\Big(\frac{p_{n-2f+i}}{p_i}\Big)^{\dim(\mu^{(i)})} \cdot \prod_{i=1}^{f-1}\Big(\frac{p_{n-2f+i+2}}{p_i}\Big)^{\dim(\mu^{(i+1)})} \cdot \Big(\frac{p_{n-2f+2}}{p_{n-2f+1}}\Big)^{\dim(\mu^{(1)})} \\
&= \prod_{i=3}^{f}\Big(\frac{p_{n-2f+i}}{p_i}\Big)^{\dim(\mu^{(i)})} \cdot \prod_{i=1}^{f-2}\Big(\frac{p_{n-2f+i+2}}{p_i}\Big)^{\dim(\mu^{(i+1)})} \\
&\qquad \times \Big(\frac{p_{n-2f+2}}{p_1}\Big)^{\dim(\mu^{(1)})+\dim(\mu^{(2)})} \Big(\frac{p_{n-f+1}}{p_f}\Big)^{\dim(\mu^{(f)})} \Big(\frac{1}{p_2}\Big)^{\dim(\mu^{(2)})} \\
&= \Big(\frac{p_{n-f+1}}{p_f}\Big)^{\dim(\mu^{(f)})} \cdot \prod_{i=1}^{f-1}\Big(\frac{p_{n-2f+i+1}}{p_i}\Big)^{\dim(\mu^{(i)})+\dim(\mu^{(i+1)})} \\
&= \prod_{i=1}^{f}\Big(\frac{p_{n-2f+i+1}}{p_i}\Big)^{\dim(\lambda^{(i)})},
\end{aligned}$$

where

$$\dim(\lambda^{(i)}) = \begin{cases} \dim\big(\mu^{(i)}\big), & \text{if } i = f; \\ \dim\big(\mu^{(i)}\big) + \dim\big(\mu^{(i+1)}\big), & \text{otherwise.} \end{cases}$$

On the other hand,

$$g_{\lambda,\lambda^{(i)}} = \Big(\frac{p_{n-2f+i+1}}{p_i}\Big)^{\dim(\lambda^{(i)})}, \qquad \text{for } i = 1, \dots, f,$$

implies that

$$\prod_{i=1}^{f} g_{\lambda,\lambda^{(i)}} = \prod_{i=1}^{f}\Big(\frac{p_{n-2f+i+1}}{p_i}\Big)^{\dim(\lambda^{(i)})}.$$

Now suppose that $n - 2f = 1$ and, for $i = 1, \dots, f-1$, let $\nu^{(i)} = (f-i-1, n+2i-2)$. Then, by induction,

$$\det\big(\nu^{(0)}\big) = \prod_{i=1}^{f-1}\left(\frac{p_{n-2f+i+1}}{p_i}\right)^{\dim(\nu^{(i)})} \qquad \text{and} \qquad \det\big(\mu^{(0)}\big) = \det\big(\nu^{(0)}\big) \cdot (p_2)^{\dim(\mu^{(0)})}.$$

Further, Lemma 5.1 and induction imply that

$$\begin{aligned}
\det(\lambda) &= \det\big(\mu^{(0)}\big) \det\big(\mu^{(1)}\big) \left(\frac{p_3}{p_2}\right)^{\dim(\mu^{(1)})} \\
&= \prod_{i=1}^{f-1}\left(\frac{p_{n-2f+i+1}}{p_i}\right)^{\dim(\nu^{(i)})} \cdot \prod_{i=1}^{f-1}\left(\frac{p_{n-2f+i+2}}{p_i}\right)^{\dim(\mu^{(i+1)})} \left(\frac{1}{p_2}\right)^{\dim(\nu^{(1)})} (p_3)^{\dim(\mu^{(1)})} \\
&= \prod_{i=1}^{f-1}\left(\frac{p_{i+2}}{p_i}\right)^{\dim(\nu^{(i)})} \cdot \prod_{i=1}^{f-1}\left(\frac{p_{i+3}}{p_i}\right)^{\dim(\mu^{(i+1)})} \left(\frac{1}{p_2}\right)^{\dim(\nu^{(1)})} (p_3)^{\dim(\mu^{(1)})}.
\end{aligned}$$

In the above expression, the exponent of p_i is j_i, where

$$j_i = \begin{cases} -\dim\bigl(\nu^{(2)}\bigr) - \dim\bigl(\mu^{(3)}\bigr) - \dim\bigl(\nu^{(1)}\bigr), & \text{if } i = 2; \\ \dim\bigl(\nu^{(1)}\bigr) - \dim\bigl(\nu^{(3)}\bigr) - \dim\bigl(\mu^{(4)}\bigr) + \dim\bigl(\mu^{(1)}\bigr), & \text{if } i = 3; \\ \dim\bigl(\nu^{(i-2)}\bigr) - \dim\bigl(\nu^{(i)}\bigr) + \dim\bigl(\mu^{(i-2)}\bigr) - \dim\bigl(\nu^{(i+1)}\bigr), & \text{if } 4 \le i < f; \\ \dim\bigl(\nu^{(i-2)}\bigr) + \dim\bigl(\mu^{(i-2)}\bigr), & \text{if } f \le i \le f+1; \\ \dim\bigl(\mu^{(i-2)}\bigr), & \text{if } i = f+2; \\ 0, & \text{otherwise.} \end{cases}$$

On the other hand, with $n - 2f = 1$,

$$\prod_{i=1}^{f} g_{\lambda,\lambda^{(i)}} = \prod_{i=1}^{f} \Bigl(\frac{p_{n-2f+i+1}}{p_i}\Bigr)^{\dim(\lambda^{(i)})} = \prod_{i=1}^{f} \Bigl(\frac{p_{i+2}}{p_i}\Bigr)^{\dim(\lambda^{(i)})}.$$

Since relative positions on the Bratteli diagram associated with $\mathcal{A}_n(x)$ imply that

$$j_i = \begin{cases} -\dim\bigl(\lambda^{(i)}\bigr), & \text{if } i = 2; \\ \dim\bigl(\lambda^{(i-2)}\bigr) - \dim\bigl(\lambda^{(i)}\bigr), & \text{if } 3 \le i \le f; \\ \dim\bigl(\lambda^{(i-2)}\bigr), & \text{if } f < i \le f+2, \end{cases}$$

the lemma holds in the case where $n-2f = 1$. If $n-2f = 0$, the given formula for $\det(\lambda)$ follows directly from Lemma 5.1. □

Example 5.1. If $n = 11$ and $\lambda = (5, 1)$, then

$$\det(\lambda) = \Bigl(\frac{p_7}{p_5}\Bigr)\Bigl(\frac{p_6}{p_4}\Bigr)^{10}\Bigl(\frac{p_5}{p_3}\Bigr)^{44}\Bigl(\frac{p_4}{p_2}\Bigr)^{110}\Bigl(\frac{p_3}{p_1}\Bigr)^{165}.$$

To write the above expression as a product in $\mathbb{Z}[x]$, we apply (3.5) with $k = 1$,

$$\begin{aligned} \det(\lambda) &= p_7\Bigl(\frac{p_3(p_4 - p_2)}{p_2(p_3 - p_1)}\Bigr)^{10}(p_5)^{43}(p_3 - p_1)^{110}(p_3)^{121} \\ &= p_7(p_3 - 2p_1)^{10}(p_5)^{43}(p_3 - p_1)^{100}(p_3)^{131}. \end{aligned}$$

Remark 5.1. The above results show that the the Temperley–Lieb algebras, besides being cellular in the sense of [2], are equipped with a family of Jucys–Murphy elements satisfying the "separation condition" defined by A. Mathas [8]. In a forthcoming note, we demonstrate a similar construction for the partition algebras of [6].

REFERENCES

[1] R. Dipper, G. James, and E. Murphy. Gram determinants of type B_n. *J. Algebra*, 189:481–505, 1997.

[2] J. Graham and G. Lehrer. Cellular algebras. *Invent. Math.*, 123(1):1–34, 1996.

[3] J. Graham and G. Lehrer. The representation theory of affine Temperley–Lieb algebras. *Enseign. Math.*, 44:173–218, 1998.

[4] T. Halverson, M. Mazzocco, and A. Ram. Commuting families in Hecke and Temperley–Lieb algebras. *Preprint*, 2007.

[5] T. Halverson and A. Ram. Characters of algebras containing a Jones Basic Construction: the Temperly-Lieb, Okada, Brauer, and Birman-Wenzl Algebras. *Adv. Math.*, 116(2):263–321, 1995.

[6] T. Halverson and A. Ram. Partition algebras. *European J. Combin.*, 26:869–921, 2005.

[7] M. Härterich. Murphy bases of generalized Temperley–Lieb algebras. *Arch. Math.*, 72:337–345, 1999.

[8] A Mathas. Seminormal forms and gram determinants for cellular algebras. *J. Reine Angew. Math.*, to appear.

[9] N. Temperley and E. Lieb. Relations between the “percolation” and “colouring” problem and other graph theoretical problems associated with regular plane lattices: Some exact results for the percolation problem. *Proc. Roy. Soc. London Ser. A*, 332:251–280, 1971.